

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

BOX PCT/DO-EO

Commissioner for Patents
Washington, D.C. 20231

"Express Mail" mailing label number **EJ 691 689 748 US**

I hereby certify that this paper is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" under 37 CFR § 1.10 on the date indicated above and addressed to: Commissioner of Patents, Washington, DC 20231, on December 1, 2000.


Colleen Hanagan

NATIONAL STAGE APPLICATION TRANSMITTAL LETTER
APPLICATION FILING UNDER 35 U.S.C. § 371

Transmitted herewith for filing is the patent application of:

Inventor(s)/Applicant(s): Vinals-Bassols, Carlota
International Application No.: PCT/EP99/03822
International Published Appln. No.: WO 99/63093
International Filing Date: 31 May 1999
Priority Filing Dates: 03 June 1998 & 08 March 1999
Thirty Month Date: 03 December 2000
Title: "BASB027 PROTEINS AND GENES FROM
MORAXELLA CATARRHALIS, ANTIGENS,
ANTIBODIES, AND USES"

1. **THIS NEW APPLICATION IS A NATIONAL STAGE APPLICATION UNDER PCT, CHAPTER II WITH A REQUEST FOR EXAMINATION WITHOUT DELAY TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US).**

- ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. § 371;
☐ This is a **SECOND** or subsequent submission of items concerning a filing under 35 U.S.C. § 371.

2. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).

3. A proper Demand for International Preliminary Examination was made by the 19TH month from the earliest claimed priority date.

4. Enclosed items are required for filing under 37 CFR § 1.53(b) and § 1.494(b) or § 1.495(b):

- ☒ One copy of International Publication **WO 99/63093**
- (a) ☒ is transmitted herewith (**required only if not transmitted by the International Bureau**)
- (b) ☐ has been transmitted by the International Bureau
- (c) ☐ is not required, as the application was filed in the United States Receiving Office (RO/US)

Fees

☒ The basic national fee set forth in 37 CFR § 1.482 - International Preliminary Examination Fee not paid to USPTO but International Search Report prepared by the EPO or JPO - **\$860.00**

- ☒ Claims in Excess of 20 (14 @ \$18.00)
- ☒ Independent Claims in Excess of 3 (2 @ \$80.00)

5. Further enclosed are:

- ☒ One copy of International Preliminary Examination Report.
- ☒ One copy of International Search Report.
- ☐ One copy of Written Opinion.
- ☐ One copy of PCT Request as filed.
- ☐ One copy of Chapter II Demand as filed.

6. ☐ A translation of the International Application into English (35 U.S.C. § 371(c)(2))

7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. § 371(c)(3))

8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. § 371(c)(3))

9. Still additional papers enclosed:

- ☐ Assignment with Assignment Recordation Form Cover Sheet
- ☐ Verification Statement Claiming Small Entity Status
- ☐ Declaration or oath is enclosed executed by the inventor
- ☐ An Information Disclosure Statement under 37 CFR § 1.97 and § 1.98
- ☒ Return Acknowledgment Postcard

Preliminary Amendment

Prior to calculation of fees, kindly enter:

- ☒ Preliminary Amendment submitted herewith
- ☐ do not enter Preliminary Amendment


10. Fee payment being made at this time is enclosed:

* Basic filing fee (\$860.00)	860.00
* Claims in Excess of 20	252.00
* (14 @ \$18.00)	
Independent Claims in Excess of 3	160.00
(2 @ \$80.00)	
* Total Fees enclosed:	<u>\$1272.00</u>

11. The Commissioner is hereby authorized to charge any fees which may be required, or credit any overpayment to Account No. 50-0258. This letter is filed in duplicate for accounting purposes.

Respectfully submitted,

Date: November 30, 2000


Arthur E. Jackson
Registration No. 34,354

DECHERT
4000 Bell Atlantic Tower
1717 Arch Street
Philadelphia, PA 19103-2793
Fax: (609) 620-3259
Attn: Arthur E. Jackson, Esq.
(609) 620-3254

101758.1.07

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Vinals-Bassols

Serial No.: Unknown

Filed: Herewith

Group Art Unit No.: Unknown

Examiner: Unknown

For: BASB027 Proteins and Genes from *Moraxella Catarrhalis*, Antigens, Antibodies, and Uses

PRELIMINARY AMENDMENT

Sir:

Please enter this Preliminary Amendment into the record of the above-identified patent application before the calculation of fees.

In the Claims:

Please delete the claims of the application as filed in the PCT and substitute therefor:

27. An isolated polypeptide comprising a member selected from the group consisting of
- (a) an amino acid sequence which has at least 90% identity to SEQ ID NOs:2 or 4;
 - (b) an immunogenic fragment of the amino acid sequence of (a), wherein the immunogenic fragment is at least 90% identical to an aligned contiguous segment of SEQ ID NOs:2 or 4; and
 - (c) an immunogenic fragment of the amino acid sequence of (a) that matches an aligned contiguous segment of SEQ ID NOs:2 or 4 with no more than five single amino acid substitutions, deletions or additions;

wherein the isolated polypeptide, when administered to a subject in a suitable composition which can include an adjuvant, or a suitable carrier coupled to the polypeptide, induces an immune response that recognizes a polypeptide having the sequence of SEQ ID NOs:2 or 4.

28. The isolated polypeptide of claim 27, wherein the immune response recognizes an antigen having molecular weight 95 kD (by SDS PAGE) in *Moraxella catarrhalis* strains Mc 2904, 2913, 2926, 2931, 2956, 2960, 2969, and 2975.

29. The isolated polypeptide of claim 27, wherein the immunogenic fragment of (b) comprises at least 15 amino acids.

30. The isolated polypeptide of claim 29, wherein the immunogenic fragment matches an aligned contiguous segment of SEQ ID NOs:2 or 4 with no more than a single amino acid substitution, deletion or addition.

31. The isolated polypeptide of claim 27, wherein the immunogenic fragment of (b) comprises at least 20 amino acids.

32. The isolated polypeptide of Claim 27 wherein the amino acid sequence of (a) has at least 95% identity to SEQ ID NOs:2 or 4.

33. The isolated polypeptide of Claim 32 wherein the isolated polypeptide comprises the amino acid sequence of SEQ ID NOs:2 or 4.

34. The isolated polypeptide of claim 32 wherein the isolated polypeptide consists of the amino acid sequence of SEQ ID NOs:2 or 4.

35. A fusion protein comprising the isolated polypeptide of Claim 27.

36. The isolated polypeptide of Claim 27 wherein the polypeptide is the immunogenic fragment having no more than two single amino acid substitutions, deletions or additions relative to the aligned sequence.

37. The isolated polypeptide of Claim 27 wherein the polypeptide is the immunogenic fragment having no more than one single amino acid substitution, deletion or addition relative to the aligned sequence.

38. The isolated polypeptide of Claim 27 wherein the polypeptide is the immunogenic fragment which matches the aligned sequence.

39. An isolated polypeptide encoded by an isolated first polynucleotide wherein the isolated first polynucleotide hybridizes under stringent conditions to a second polynucleotide which encodes the polypeptide of SEQ ID NOs:2 or 4; wherein stringent conditions comprise overnight incubation at 42° C in a solution comprising: 50% formamide, 5×SSC (150 mM

NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5× Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1× SSC at about 65° C; wherein the isolated polypeptide, when administered to a subject in a suitable composition which can include an adjuvant, or a suitable carrier coupled to the polypeptide, induces an immune response that recognizes a polypeptide having the sequence of SEQ ID NOs:2 or 4.

40. An isolated polynucleotide encoding a polypeptide of Claim 27 or the full complement to the isolated polynucleotide.

41. An isolated polynucleotide encoding a polypeptide of Claim 27, wherein the isolated polynucleotide encodes the polypeptide comprising SEQ ID NOs:2 or 4.

42. An isolated polynucleotide comprising the polynucleotide of SEQ ID NOs:1 or 3.

43. An isolated polynucleotide segment comprising a polynucleotide sequence or the full complement of the entire length of the polynucleotide sequence, wherein the polynucleotide sequence hybridizes to the full complement of SEQ ID NOs:1 or 3 minus the complement of any stop codon, wherein the hybridization conditions include incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml denatured, sheared salmon sperm DNA, followed by washing in 0.1x SSC at 65°C; and, wherein the polynucleotide sequence is identical to SEQ ID NOs:1 or 3 minus any terminal stop codon, except that, over the entire length corresponding to SEQ ID NOs:1 or 3 minus any terminal stop codon, n_n nucleotides are substituted, inserted or deleted, wherein n_n satisfies the following expression

$$n_n \leq x_n - (x_n \cdot y)$$

wherein x_n is the total number of nucleotides in SEQ ID NOs:1 or 3 minus any terminal stop codon, y is at least 0.95, and wherein any non-integer product of x_n and y is rounded down to the nearest integer before subtracting the product from x_n ; and wherein the polynucleotide sequence detects *Moraxella catarrhalis*.

44. An expression vector comprising the isolated polynucleotide of Claim 40.

45. A host cell transformed with the expression vector of Claim 44.
46. A process of producing an isolated polypeptide comprising (a) culturing the host cell of Claim 45 under conditions sufficient for the production of the encoded polypeptide and (b) recovering the polypeptide.
47. A nucleic acid vaccine comprising the isolated polynucleotide of Claim 40 and a pharmaceutically acceptable carrier.
48. An isolated polynucleotide segment comprising a polynucleotide sequence or the full complement of the entire length of the polynucleotide sequence, wherein the polynucleotide sequence is identical to SEQ ID NOs:1 or 3 minus any terminal stop codon, except that, over the entire length corresponding to SEQ ID NOs:1 or 3 minus any terminal stop codon, n_n nucleotides are substituted, inserted or deleted, wherein n_n satisfies the following expression
- $$n_n \leq x_n - (x_n \bullet y)$$
- wherein x_n is the total number of nucleotides in SEQ ID NOs:1 or 3 minus any terminal stop codon, y is at least 0.90, and wherein any non-integer product of x_n and y is rounded down to the nearest integer before subtracting the product from x_n ; and wherein the polynucleotide sequence detects *Moraxella catarrhalis*.
49. The isolated polynucleotide of Claim 48 where y is at least 0.95.
50. An expression vector comprising the isolated polynucleotide of Claim 48 which codes for a polypeptide that, when administered to a mammal which can include an adjuvant, or a suitable carrier coupled to the polypeptide, induces an immune response that recognizes a polypeptide having the sequence of SEQ ID NOs:2 or 4.
51. A host cell transformed with the isolated polynucleotide or an expression vector comprising the isolated polynucleotide of Claim 48.

52. A process of producing an isolated polypeptide comprising (a) culturing the host cell of Claim 51 under conditions sufficient for the production of the encoded polypeptide and (b) recovering the polypeptide.

53. A vaccine comprising the polypeptide of Claim 27 and a pharmaceutically acceptable carrier.

54. The vaccine of Claim 53, wherein the composition comprises at least one other *Moraxella catarrhalis* antigen.

55. An antibody immunospecific for the polypeptide or immunogenic fragment of Claim 27.

56. A method for inducing an immune response in a mammal comprising administration of the polypeptide of Claim 27.

57. A method of diagnosing a *Moraxella catarrhalis* infection, comprising identifying a polypeptide of Claim 27, or an antibody that is immunospecific for the polypeptide, present within a biological sample from an animal suspected of having such an infection.

58. A method for inducing an immune response in a mammal comprising administration of the isolated polynucleotide of Claim 40.

59. A therapeutic composition useful in treating humans with *Moraxella catarrhalis* comprising at least one antibody directed against the polypeptide of claim 27 and a suitable pharmaceutical carrier.

60. A process for expressing the polynucleotide of Claim 40 comprising transforming a host cell with the expression vector comprising the polynucleotide and culturing the host cell under conditions sufficient for expression of the polynucleotide.

REMARKS

Applicant respectfully requests that this Preliminary Amendment be entered in this case before the calculation of fees and before examination of the subject application.

Claims

Claims 1-26 have been canceled without prejudice or disclaimer of the subject matter therein. Applicant reserves the right to prosecute, in one or more patent applications, the canceled claims, the claims to non-elected inventions, the claims as originally filed, and any other claims supported by the specification.

New claims 27-60 have been introduced. No new matter is added.

Support

Support for the new claims is either obvious, or is as described in the text below. Particularly, support for the recitation of "five single amino acid substitutions, deletions or additions" may be found, for example, at page 7, lines 1-2. Support for compositions of the isolated polypeptide which include an adjuvant recited in the claims may be found, for example, at page 40, lines 28-29. Support for the stringent hybridization conditions may be found, for example, at page 14, lines 20-24. Support for the recitation of sequence relatedness such as "wherein the first polynucleotide sequence is identical to SEQ ID NOs:1 or 3, except that, over the entire length corresponding to SEQ ID NOs:1 or 3, n_n nucleotides are substituted, inserted or deleted, wherein n_n satisfies the following expression

$$n_n \leq x_n - (x_n \bullet y)$$


wherein x_n is the total number of nucleotides in SEQ ID NOs:1 or 3, y is at least 0.90, and wherein any non-integer product of x_n and y is rounded down to the nearest integer before subtracting the product from x_n " may be found in the specification, for example, at page 44, line 7 through page 45, line 2. Support for the recitation, "wherein the immune response recognizes an antigen having molecular weight 95 kD (by SDS PAGE) in *Moraxella catarrhalis* strains Mc 2931, 2904, 2905, 2906, 2907, 2908, 2909, 2910 and 2911" can be found, for example, in Example 6 at page 63, lines 4 and 5 and Figure 7.

International Application No.: PCT/EP99/03822
Attorney Docket No.: BM45324

Closing Remarks

Entry of this Preliminary Amendment prior to calculation of fees and Examination and Allowance of the pending claims are respectfully requested

Respectfully submitted,



Arthur E. Jackson
Registration No. 34,354

DECHERT PRICE & RHOADS
Princeton Pike Corporate Center
PO Box 5218
Princeton, New Jersey 08543-5218
Arthur E. Jackson (609) 620-3254
Fax: (609) 620-3259
Attn: Arthur E. Jackson, Esq.
(609) 620-3254

FIELD OF THE INVENTION

This invention relates to polynucleotides, (herein referred to as "BASB027 polynucleotide(s)"), polypeptides encoded by them (referred to herein as "BASB027" or "BASB027 polypeptide(s)"), recombinant materials and methods for their production. In another aspect, the invention relates to methods for using such polypeptides and polynucleotides, including vaccines against bacterial infections. In a further aspect, the invention relates to diagnostic assays for detecting infection of certain pathogens.

BACKGROUND OF THE INVENTION

Moraxella catarrhalis (also named *Branhamella catarrhalis*) is a Gram negative bacteria frequently isolated from the human upper respiratory tract. It is responsible for several pathologies the main ones being otitis media in infants and children, and pneumonia in elderlies. It is also responsible of sinusitis, nosocomial infections and less frequently of invasive diseases.

Otitis media is an important childhood disease both by the number of cases and its potential sequelae. More than 3.5 millions cases are recorded every year in the United States, and it is estimated that 80 % of the children have experienced at least one episode of otitis before reaching the age of 3 (Klein, JO (1994) Clin.Inf.Dis 19:823). Left untreated, or becoming chronic, this disease may lead to hearing losses that could be temporary (in the case of fluid accumulation in the middle ear) or permanent (if the auditive nerve is damaged). In infants, such hearing losses may be responsible for a delayed speech learning.

Three bacterial species are primarily isolated from the middle ear of children with otitis media: *Streptococcus pneumoniae*, non typeable *Haemophilus influenza* (NTHi) and *M. catarrhalis*. They are present in 60 to 90 % of the cases. A review of recent studies shows that *S. pneumoniae* and NTHi represent both about 30 %, and *M. catarrhalis* about 15 % of

the otitis media cases (Murphy, TF (1996) Microbiol.Rev. 60:267). Other bacteria could be isolated from the middle ear (*H influenza* type B, *S. pyogenes* etc) but at a much lower frequency (2 % of the cases or less).

Epidemiological data indicate that, for the pathogens found in the middle ear, the colonization of the upper respiratory tract is an absolute prerequisite for the development of an otitis; other are however also required to lead to the disease (Dickinson, DP et al. (1988) J. Infect.Dis. 158:205, Faden, HL et al. (1991) Ann.Otorhinol.Laryngol. 100:612). These are important to trigger the migration of the bacteria into the middle ear via the Eustachian tubes, followed by the initiation of an inflammatory process. These factors are unknown todate. It has been postulated that a transient anomaly of the immune system following a viral infection, for example, could cause an inability to control the colonization of the respiratory tract (Faden, HL et al (1994) J. Infect.Dis. 169:1312). An alternative explanation is that the exposure to environmental factors allow a more important colonization of some children, who subsequently become susceptible to the development of otitis media because of the sustained presence of middle ear pathogens (Murphy, TF (1996) Microbiol.Rev. 60:267).

The immune response to *M. catarrhalis* is poorly characterized. The analysis of strains isolated sequentially from the nasopharynx of babies followed from 0 to 2 years of age, indicates that they get and eliminate frequently new strains. This indicates that an efficacious immune response against this bacteria is mounted by the colonized children (Faden, HL et al (1994) J. Infect.Dis. 169:1312).

In most adults tested, bactericidal antibodies have been identified (Chapman, AJ et al. (1985) J. Infect.Dis. 151:878). Strains of *M. catarrhalis* present variations in their capacity to resist serum bactericidal activity: in general, isolates from diseased individuals are more resistant than those who are simply colonized (Hol, C et al. (1993) Lancet 341:1281, Jordan, KL et al. (1990) Am.J.Med. 88 (suppl. 5A):28S). Serum resistance could therefore be

considered as a virulence factor of the bacteria. An opsonizing activity has been observed in the sera of children recovering from otitis media.

The antigens targetted by these different immune responses in humans have not been identified, with the exception of OMP B1, a 84 kDa protein which expression is regulated by iron, and that is recognized by the sera of patients with pneumonia (Sethi, S. et al. (1995) Infect.Immun. 63:1516), and of UspA1 and UspA2 (Chen D. et al.(1999), Infect.Immun. 67:1310).

A few other membrane proteins present on the surface of *M. catarrhalis* have been characterized using biochemical method, or for their potential implication in the induction of a protective immunity (for review, see Murphy, TF (1996) Microbiol.Rev. 60:267). In a mouse pneumonia model, the presence of antibodies raised against some of them (UspA, CopB) favors a faster clearance of the pulmonary infection. Another polypeptide (OMP CD) is highly conserved among *M. catarrhalis* strains, and presents homologies with a porin of *Pseudomonas aeruginosa*, which has been demonstrated efficacious against this bacterium in animal models.

The frequency of *Moraxella catarrhalis* infections has risen dramatically in the past few decades. This has been attributed to the emergence of multiply antibiotic resistant strains and an increasing population of people with weakened immune systems. It is no longer uncommon to isolate *Moraxella catarrhalis* strains that are resistant to some or all of the standard antibiotics. This phenomenon has created an unmet medical need and demand for new anti-microbial agents, vaccines, drug screening methods, and diagnostic tests for this organism.

SUMMARY OF THE INVENTION

The present invention relates to BASB027, in particular BASB027 polypeptides and BASB027 polynucleotides, recombinant materials and methods for their production. In another aspect, the invention relates to methods for using such polypeptides and polynucleotides, including prevention and treatment of microbial diseases, amongst others. In a further aspect, the invention relates to diagnostic assays for detecting diseases associated with microbial infections and conditions associated with such infections, such as assays for detecting expression or activity of BASB027 polynucleotides or polypeptides.

Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following descriptions and from reading the other parts of the present disclosure.

DESCRIPTION OF THE INVENTION

The invention relates to BASB027 polypeptides and polynucleotides as described in greater detail below. In particular, the invention relates to polypeptides and polynucleotides of BASB027 of *Moraxella catarrhalis*, which is related by amino acid sequence homology to *Neisseria meningitidis* OMP85 outer membrane protein. The invention relates especially to BASB027 having the nucleotide and amino acid sequences set out in SEQ ID NO:1 or 3 and SEQ ID NO:2 or 4 respectively. It is understood that sequences recited in the Sequence Listing below as "DNA" represent an exemplification of one embodiment of the invention, since those of ordinary skill will recognize that such sequences can be usefully employed in polynucleotides in general, including ribopolynucleotides.

Polypeptides

In one aspect of the invention there are provided polypeptides of *Moraxella catarrhalis* referred to herein as "BASB027" and "BASB027 polypeptides" as well as biologically, diagnostically, prophylactically, clinically or therapeutically useful variants thereof, and compositions comprising the same.

The present invention further provides for:

- (a) an isolated polypeptide which comprises an amino acid sequence which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, most preferably at least 97-99% or exact identity, to that of SEQ ID NO:2 or 4;
- (b) a polypeptide encoded by an isolated polynucleotide comprising a polynucleotide sequence which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, even more preferably at least 97-99% or exact identity to SEQ ID NO:1 or 3 over the entire length of SEQ ID NO:1 or 3 respectively; or
- (c) a polypeptide encoded by an isolated polynucleotide comprising a polynucleotide sequence encoding a polypeptide which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, even more preferably at least 97-99% or exact identity, to the amino acid sequence of SEQ ID NO:2 or 4.

The BASB027 polypeptides provided in SEQ ID NO:2 or 4 are the BASB027 polypeptides from *Moraxella catarrhalis* strain Mc2931 (ATCC 43617).

The invention also provides an immunogenic fragment of a BASB027 polypeptide, that is, a contiguous portion of the BASB027 polypeptide which has the same or substantially the same immunogenic activity as the polypeptide comprising the amino acid sequence of SEQ ID NO:2 or 4; That is to say, the fragment (if necessary when coupled to a carrier) is capable of raising an immune response which recognises the BASB027 polypeptide.

Such an immunogenic fragment may include, for example, the BASB027 polypeptide lacking an N-terminal leader sequence, and/or a transmembrane domain and/or a C-terminal anchor domain. In a preferred aspect the immunogenic fragment of BASB027 according to the invention comprises substantially all of the extracellular domain of a polypeptide which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, most preferably at least 97-99% identity, to that of SEQ ID NO:2 or 4 over the entire length of SEQ ID NO:2

A fragment is a polypeptide having an amino acid sequence that is entirely the same as part but not all of any amino acid sequence of any polypeptide of the invention. As with BASB027 polypeptides, fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most preferably as a single continuous region in a single larger polypeptide.

Preferred fragments include, for example, truncation polypeptides having a portion of an amino acid sequence of SEQ ID NO:2 or 4 or of variants thereof, such as a continuous series of residues that includes an amino- and/or carboxyl-terminal amino acid sequence. Degradation forms of the polypeptides of the invention produced by or in a host cell, are also preferred. Further preferred are fragments characterized by structural or functional attributes such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions.

Further preferred fragments include an isolated polypeptide comprising an amino acid sequence having at least 15, 20, 30, 40, 50 or 100 contiguous amino acids from the amino acid sequence of SEQ ID NO:2 or 4, or an isolated polypeptide comprising an amino acid sequence having at least 15, 20, 30, 40, 50 or 100 contiguous amino acids truncated or deleted from the amino acid sequence of SEQ ID NO:2 or 4.

Fragments of the polypeptides of the invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, these fragments may be employed as intermediates for producing the full-length polypeptides of the invention.

Particularly preferred are variants in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acids are substituted, deleted, or added in any combination.

The polypeptides, or immunogenic fragments, of the invention may be in the form of the "mature" protein or may be a part of a larger protein such as a precursor or a fusion protein. It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production. Furthermore, addition of exogenous polypeptide or lipid tail or polynucleotide sequences to increase the immunogenic potential of the final molecule is also considered.

In one aspect, the invention relates to genetically engineered soluble fusion proteins comprising a polypeptide of the present invention, or a fragment thereof, and various portions of the constant regions of heavy or light chains of immunoglobulins of various subclasses (IgG, IgM, IgA, IgE). Preferred as an immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG1, where fusion takes place at the hinge region. In a particular embodiment, the Fc part can be removed simply by incorporation of a cleavage sequence which can be cleaved with blood clotting factor Xa.

Furthermore, this invention relates to processes for the preparation of these fusion proteins by genetic engineering, and to the use thereof for drug screening, diagnosis and therapy. A further aspect of the invention also relates to polynucleotides encoding such fusion proteins. Examples of fusion protein technology can be found in International Patent Application Nos. WO94/29458 and WO94/22914.

The proteins may be chemically conjugated, or expressed as recombinant fusion proteins allowing increased levels to be produced in an expression system as compared to non-fused protein. The fusion partner may assist in providing T helper epitopes

(immunological fusion partner), preferably T helper epitopes recognised by humans, or assist in expressing the protein (expression enhancer) at higher yields than the native recombinant protein. Preferably the fusion partner will be both an immunological fusion partner and expression enhancing partner.

Fusion partners include protein D from *Haemophilus influenzae* and the non-structural protein from influenzae virus, NS1 (hemagglutinin). Another fusion partner is the protein known as LytA. Preferably the C terminal portion of the molecule is used. LytA is derived from *Streptococcus pneumoniae* which synthesize an N-acetyl-L-alanine amidase, amidase LytA, (coded by the lytA gene {Gene, 43 (1986) page 265-272}) an autolysin that specifically degrades certain bonds in the peptidoglycan backbone. The C-terminal domain of the LytA protein is responsible for the affinity to the choline or to some choline analogues such as DEAE. This property has been exploited for the development of E.coli C-LytA expressing plasmids useful for expression of fusion proteins. Purification of hybrid proteins containing the C-LytA fragment at its amino terminus has been described {Biotechnology: 10, (1992) page 795-798}. It is possible to use the repeat portion of the LytA molecule found in the C terminal end starting at residue 178, for example residues 188 - 305.

The present invention also includes variants of the aforementioned polypeptides, that is polypeptides that vary from the referents by conservative amino acid substitutions, whereby a residue is substituted by another with like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr.

Polypeptides of the present invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a

combination of these methods. Means for preparing such polypeptides are well understood in the art.

It is most preferred that a polypeptide of the invention is derived from *Moraxella catarrhalis*, however, it may preferably be obtained from other organisms of the same taxonomic genus. A polypeptide of the invention may also be obtained, for example, from organisms of the same taxonomic family or order.

Polynucleotides

It is an object of the invention to provide polynucleotides that encode BASB027 polypeptides, particularly polynucleotides that encode the polypeptide herein designated BASB027.

In a particularly preferred embodiment of the invention the polynucleotide comprises a region encoding BASB027 polypeptides comprising a sequence set out in SEQ ID NO:1 or 3 which includes a full length gene, or a variant thereof.

The BASB027 polynucleotides provided in SEQ ID NO:1 or 3 are the BASB027 polynucleotides from *Moraxella catarrhalis* strain Mc2931 (ATCC 43617).

As a further aspect of the invention there are provided isolated nucleic acid molecules encoding and/or expressing BASB027 polypeptides and polynucleotides, particularly *Moraxella catarrhalis* BASB027 polypeptides and polynucleotides, including, for example, unprocessed RNAs, ribozyme RNAs, mRNAs, cDNAs, genomic DNAs, B- and Z-DNAs. Further embodiments of the invention include biologically, diagnostically, prophylactically, clinically or therapeutically useful polynucleotides and polypeptides, and variants thereof, and compositions comprising the same.

Another aspect of the invention relates to isolated polynucleotides, including at least one full length gene, that encodes a BASB027 polypeptide having a deduced amino acid sequence of SEQ ID NO:2 or 4 and polynucleotides closely related thereto and variants thereof.

In another particularly preferred embodiment of the invention there is a BASB027 polypeptide from *Moraxella catarrhalis* comprising or consisting of an amino acid sequence of SEQ ID NO:2 or 4 or a variant thereof.

Using the information provided herein, such as a polynucleotide sequence set out in SEQ ID NO:1 or 3, a polynucleotide of the invention encoding BASB027 polypeptide may be obtained using standard cloning and screening methods, such as those for cloning and sequencing chromosomal DNA fragments from bacteria using *Moraxella catarrhalis* Catlin cells as starting material, followed by obtaining a full length clone. For example, to obtain a polynucleotide sequence of the invention, such as a polynucleotide sequence given in SEQ ID NO:1 or 3, typically a library of clones of chromosomal DNA of *Moraxella catarrhalis* Catlin in *E.coli* or some other suitable host is probed with a radiolabeled oligonucleotide, preferably a 17-mer or longer, derived from a partial sequence. Clones carrying DNA identical to that of the probe can then be distinguished using stringent hybridization conditions. By sequencing the individual clones thus identified by hybridization with sequencing primers designed from the original polypeptide or polynucleotide sequence it is then possible to extend the polynucleotide sequence in both directions to determine a full length gene sequence. Conveniently, such sequencing is performed, for example, using denatured double stranded DNA prepared from a plasmid clone. Suitable techniques are described by Maniatis, T., Fritsch, E.F. and Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989). (see in particular Screening By Hybridization 1.90 and Sequencing Denatured Double-Stranded DNA Templates 13.70). Direct genomic DNA sequencing may also be performed to obtain a full length gene

sequence. Illustrative of the invention, each polynucleotide set out in SEQ ID NO:1 or 3 was discovered in a DNA library derived from *Moraxella catarrhalis*.

Moreover, each DNA sequence set out in SEQ ID NO:1 or 3 contains an open reading frame encoding a protein having about the number of amino acid residues set forth in SEQ ID NO:2 or 4 with a deduced molecular weight that can be calculated using amino acid residue molecular weight values well known to those skilled in the art.

The polynucleotide of SEQ ID NO:1, between the start codon at nucleotide number 1 and the stop codon which begins at nucleotide number 2440 of SEQ ID NO:1, encodes the polypeptide of SEQ ID NO:2.

The polynucleotide of SEQ ID NO:3, between the start codon at nucleotide number 1 and the stop codon which begins at nucleotide number 2440 of SEQ ID NO:3, encodes the polypeptide of SEQ ID NO:4.

In a further aspect, the present invention provides for an isolated polynucleotide comprising or consisting of:

- (a) a polynucleotide sequence which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, even more preferably at least 97-99% or exact identity to SEQ ID NO:1 or 3 over the entire length of SEQ ID NO:1 or 3 respectively; or
- (b) a polynucleotide sequence encoding a polypeptide which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, even more preferably at least 97-99% or 100% exact, to the amino acid sequence of SEQ ID NO:2 or 4, over the entire length of SEQ ID NO:2 or 4 respectively.

A polynucleotide encoding a polypeptide of the present invention, including homologs and orthologs from species other than *Moraxella catarrhalis*, may be obtained by a process which comprises the steps of screening an appropriate library under stringent hybridization

conditions (for example, using a temperature in the range of 45 – 65°C and an SDS concentration from 0.1 – 1%) with a labeled or detectable probe consisting of or comprising the sequence of SEQ ID NO:1 or 3 or a fragment thereof; and isolating a full-length gene and/or genomic clones containing said polynucleotide sequence.

The invention provides a polynucleotide sequence identical over its entire length to a coding sequence (open reading frame) in SEQ ID NO:1 or 3. Also provided by the invention is a coding sequence for a mature polypeptide or a fragment thereof, by itself as well as a coding sequence for a mature polypeptide or a fragment in reading frame with another coding sequence, such as a sequence encoding a leader or secretory sequence, a pre-, or pro- or prepro-protein sequence. The polynucleotide of the invention may also contain at least one non-coding sequence, including for example, but not limited to at least one non-coding 5' and 3' sequence, such as the transcribed but non-translated sequences, termination signals (such as rho-dependent and rho-independent termination signals), ribosome binding sites, Kozak sequences, sequences that stabilize mRNA, introns, and polyadenylation signals. The polynucleotide sequence may also comprise additional coding sequence encoding additional amino acids. For example, a marker sequence that facilitates purification of the fused polypeptide can be encoded. In certain embodiments of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, *Proc. Natl. Acad. Sci., USA* 86: 821-824 (1989), or an HA peptide tag (Wilson *et al.*, *Cell* 37: 767 (1984), both of which may be useful in purifying polypeptide sequence fused to them. Polynucleotides of the invention also include, but are not limited to, polynucleotides comprising a structural gene and its naturally associated sequences that control gene expression.

The nucleotide sequence encoding BASB027 polypeptide of SEQ ID NO:2 or 4 may be identical to the polypeptide encoding sequence contained in nucleotides 1 to 2439 of SEQ ID NO:1 or 3 respectively. Alternatively it may be a sequence, which as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID

NO:2 or 4.

The term "polynucleotide encoding a polypeptide" as used herein encompasses polynucleotides that include a sequence encoding a polypeptide of the invention, particularly a bacterial polypeptide and more particularly a polypeptide of the *Moraxella catarrhalis* BASB027 having an amino acid sequence set out in SEQ ID NO:2 or 4. The term also encompasses polynucleotides that include a single continuous region or discontinuous regions encoding the polypeptide (for example, polynucleotides interrupted by integrated phage, an integrated insertion sequence, an integrated vector sequence, an integrated transposon sequence, or due to RNA editing or genomic DNA reorganization) together with additional regions, that also may contain coding and/or non-coding sequences.

The invention further relates to variants of the polynucleotides described herein that encode variants of a polypeptide having a deduced amino acid sequence of SEQ ID NO:2 or 4. Fragments of polynucleotides of the invention may be used, for example, to synthesize full-length polynucleotides of the invention.

Further particularly preferred embodiments are polynucleotides encoding BASB027 variants, that have the amino acid sequence of BASB027 polypeptide of SEQ ID NO:2 or 4 in which several, a few, 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues are substituted, modified, deleted and/or added, in any combination. Especially preferred among these are silent substitutions, additions and deletions, that do not alter the properties and activities of BASB027 polypeptide.

Further preferred embodiments of the invention are polynucleotides that are at least 85% identical over their entire length to a polynucleotide encoding BASB027 polypeptide having an amino acid sequence set out in SEQ ID NO:2 or 4, and polynucleotides that are complementary to such polynucleotides. Alternatively, most highly preferred are polynucleotides that comprise a region that is at least 90% identical over its entire length to

a polynucleotide encoding BASB027 polypeptide and polynucleotides complementary thereto. In this regard, polynucleotides at least 95% identical over their entire length to the same are particularly preferred. Furthermore, those with at least 97% are highly preferred among those with at least 95%, and among these those with at least 98% and at least 99% are particularly highly preferred, with at least 99% being the more preferred.

Preferred embodiments are polynucleotides encoding polypeptides that retain substantially the same biological function or activity as the mature polypeptide encoded by a DNA of SEQ ID NO:1 or 3.

In accordance with certain preferred embodiments of this invention there are provided polynucleotides that hybridize, particularly under stringent conditions, to BASB027 polynucleotide sequences, such as those polynucleotides in SEQ ID NO:1 or 3.

The invention further relates to polynucleotides that hybridize to the polynucleotide sequences provided herein. In this regard, the invention especially relates to polynucleotides that hybridize under stringent conditions to the polynucleotides described herein. As herein used, the terms "stringent conditions" and "stringent hybridization conditions" mean hybridization occurring only if there is at least 95% and preferably at least 97% identity between the sequences. A specific example of stringent hybridization conditions is overnight incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml of denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1x SSC at about 65°C. Hybridization and wash conditions are well known and exemplified in Sambrook, *et al.*, *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor, N.Y., (1989), particularly Chapter 11 therein. Solution hybridization may also be used with the polynucleotide sequences provided by the invention.

The invention also provides a polynucleotide consisting of or comprising a polynucleotide sequence obtained by screening an appropriate library containing the complete gene for a polynucleotide sequence set forth in SEQ ID NO:1 or 3 under stringent hybridization conditions with a probe having the sequence of said polynucleotide sequence set forth in SEQ ID NO:1 or 3 or a fragment thereof; and isolating said polynucleotide sequence. Fragments useful for obtaining such a polynucleotide include, for example, probes and primers fully described elsewhere herein.

As discussed elsewhere herein regarding polynucleotide assays of the invention, for instance, the polynucleotides of the invention, may be used as a hybridization probe for RNA, cDNA and genomic DNA to isolate full-length cDNAs and genomic clones encoding BASB027 and to isolate cDNA and genomic clones of other genes that have a high identity, particularly high sequence identity, to the BASB027 gene. Such probes generally will comprise at least 15 nucleotide residues or base pairs. Preferably, such probes will have at least 30 nucleotide residues or base pairs and may have at least 50 nucleotide residues or base pairs. Particularly preferred probes will have at least 20 nucleotide residues or base pairs and will have less than 30 nucleotide residues or base pairs.

A coding region of a BASB027 gene may be isolated by screening using a DNA sequence provided in SEQ ID NO:1 or 3 to synthesize an oligonucleotide probe. A labeled oligonucleotide having a sequence complementary to that of a gene of the invention is then used to screen a library of cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

There are several methods available and well known to those skilled in the art to obtain full-length DNAs, or extend short DNAs, for example those based on the method of Rapid Amplification of cDNA ends (RACE) (see, for example, Frohman, *et al.*, *PNAS USA* 85: 8998-9002, 1988). Recent modifications of the technique, exemplified by the Marathon™ technology (Clontech Laboratories Inc.) for example, have significantly simplified the

search for longer cDNAs. In the Marathon™ technology, cDNAs have been prepared from mRNA extracted from a chosen tissue and an 'adaptor' sequence ligated onto each end. Nucleic acid amplification (PCR) is then carried out to amplify the "missing" 5' end of the DNA using a combination of gene specific and adaptor specific oligonucleotide primers. The PCR reaction is then repeated using "nested" primers, that is, primers designed to anneal within the amplified product (typically an adaptor specific primer that anneals further 3' in the adaptor sequence and a gene specific primer that anneals further 5' in the selected gene sequence). The products of this reaction can then be analyzed by DNA sequencing and a full-length DNA constructed either by joining the product directly to the existing DNA to give a complete sequence, or carrying out a separate full-length PCR using the new sequence information for the design of the 5' primer.

The polynucleotides and polypeptides of the invention may be employed, for example, as research reagents and materials for discovery of treatments of and diagnostics for diseases, particularly human diseases, as further discussed herein relating to polynucleotide assays.

The polynucleotides of the invention that are oligonucleotides derived from a sequence of SEQ ID NOS:1 or 3 may be used in the processes herein as described, but preferably for PCR, to determine whether or not the polynucleotides identified herein in whole or in part are transcribed in bacteria in infected tissue. It is recognized that such sequences will also have utility in diagnosis of the stage of infection and type of infection the pathogen has attained.

The invention also provides polynucleotides that encode a polypeptide that is the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature polypeptide (when the mature form has more than one polypeptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, may allow protein transport, may lengthen or shorten protein half-life or may facilitate manipulation of a protein for assay or production, among other things. As

generally is the case *in vivo*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

For each and every polynucleotide of the invention there is provided a polynucleotide complementary to it. It is preferred that these complementary polynucleotides are fully complementary to each polynucleotide with which they are complementary.

A precursor protein, having a mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. When prosequences are removed such inactive precursors generally are activated. Some or all of the prosequences may be removed before activation. Generally, such precursors are called proproteins.

In addition to the standard A, G, C, T/U representations for nucleotides, the term "N" may also be used in describing certain polynucleotides of the invention. "N" means that any of the four DNA or RNA nucleotides may appear at such a designated position in the DNA or RNA sequence, except it is preferred that N is not a nucleic acid that when taken in combination with adjacent nucleotide positions, when read in the correct reading frame, would have the effect of generating a premature termination codon in such reading frame.

In sum, a polynucleotide of the invention may encode a mature protein, a mature protein plus a leader sequence (which may be referred to as a preprotein), a precursor of a mature protein having one or more prosequences that are not the leader sequences of a preprotein, or a preproprotein, which is a precursor to a proprotein, having a leader sequence and one or more prosequences, which generally are removed during processing steps that produce active and mature forms of the polypeptide.

In accordance with an aspect of the invention, there is provided the use of a polynucleotide of the invention for therapeutic or prophylactic purposes, in particular genetic immunization.

The use of a polynucleotide of the invention in genetic immunization will preferably employ a suitable delivery method such as direct injection of plasmid DNA into muscles (Wolff *et al.*, *Hum Mol Genet* (1992) 1: 363, Manthorpe *et al.*, *Hum. Gene Ther.* (1983) 4: 419), delivery of DNA complexed with specific protein carriers (Wu *et al.*, *J Biol Chem.* (1989) 264: 16985), coprecipitation of DNA with calcium phosphate (Benvenisty & Reshef, *PNAS USA*, (1986) 83: 9551), encapsulation of DNA in various forms of liposomes (Kaneda *et al.*, *Science* (1989) 243: 375), particle bombardment (Tang *et al.*, *Nature* (1992) 356:152, Eisenbraun *et al.*, *DNA Cell Biol* (1993) 12: 791) and *in vivo* infection using cloned retroviral vectors (Seeger *et al.*, *PNAS USA* (1984) 81: 5849).

Vectors, Host Cells, Expression Systems

The invention also relates to vectors that comprise a polynucleotide or polynucleotides of the invention, host cells that are genetically engineered with vectors of the invention and the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the invention.

Recombinant polypeptides of the present invention may be prepared by processes well known in those skilled in the art from genetically engineered host cells comprising expression systems. Accordingly, in a further aspect, the present invention relates to expression systems that comprise a polynucleotide or polynucleotides of the present invention, to host cells which are genetically engineered with such expression systems, and to the production of polypeptides of the invention by recombinant techniques.

For recombinant production of the polypeptides of the invention, host cells can be genetically engineered to incorporate expression systems or portions thereof or polynucleotides of the invention. Introduction of a polynucleotide into the host cell can be

effected by methods described in many standard laboratory manuals, such as Davis, *et al.*, *BASIC METHODS IN MOLECULAR BIOLOGY*, (1986) and Sambrook, *et al.*, *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989), such as, calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction and infection.

Representative examples of appropriate hosts include bacterial cells, such as cells of streptococci, staphylococci, enterococci, *E. coli*, streptomyces, cyanobacteria, *Bacillus subtilis*, *Neisseria meningitidis* and *Moraxella catarrhalis*; fungal cells, such as cells of a yeast, *Kluveromyces*, *Saccharomyces*, a basidiomycete, *Candida albicans* and *Aspergillus*; insect cells such as cells of *Drosophila* S2 and *Spodoptera* Sf9; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, 293, CV-1 and Bowes melanoma cells; and plant cells, such as cells of a gymnosperm or angiosperm.

A great variety of expression systems can be used to produce the polypeptides of the invention. Such vectors include, among others, chromosomal-, episomal- and virus-derived vectors, for example, vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses, picornaviruses, retroviruses, and alphaviruses and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression system constructs may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides and/or to express a polypeptide in a host may be used for expression in this regard. The appropriate DNA sequence may be inserted into the expression system by any

of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook *et al.*, *MOLECULAR CLONING, A LABORATORY MANUAL*, (*supra*).

In recombinant expression systems in eukaryotes, for secretion of a translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the expressed polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

Polypeptides of the present invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, ion metal affinity chromatography (IMAC) is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during intracellular synthesis, isolation and or purification.

The expression system may also be a recombinant live microorganism, such as a virus or bacterium. The gene of interest can be inserted into the genome of a live recombinant virus or bacterium. Inoculation and *in vivo* infection with this live vector will lead to *in vivo* expression of the antigen and induction of immune responses. Viruses and bacteria used for this purpose are for instance: poxviruses (e.g; vaccinia, fowlpox, canarypox), alphaviruses (Sindbis virus, Semliki Forest Virus, Venezuelan Equine Encephalitis Virus), adenoviruses, adeno-associated virus, picornaviruses (poliovirus, rhinovirus), herpesviruses (varicella zoster virus, etc), Listeria, Salmonella, Shigella, BCG. These viruses and bacteria can be virulent, or attenuated in various ways in order to obtain live vaccines. Such live vaccines also form part of the invention.

Diagnostic, Prognostic, Serotyping and Mutation Assays

This invention is also related to the use of BASB027 polynucleotides and polypeptides of the invention for use as diagnostic reagents. Detection of BASB027 polynucleotides and/or polypeptides in a eukaryote, particularly a mammal, and especially a human, will provide a diagnostic method for diagnosis of disease, staging of disease or response of an infectious organism to drugs. Eukaryotes, particularly mammals, and especially humans, particularly those infected or suspected to be infected with an organism comprising the BASB027 gene or protein, may be detected at the nucleic acid or amino acid level by a variety of well known techniques as well as by methods provided herein.

Polypeptides and polynucleotides for prognosis, diagnosis or other analysis may be obtained from a putatively infected and/or infected individual's bodily materials. Polynucleotides from any of these sources, particularly DNA or RNA, may be used directly for detection or may be amplified enzymatically by using PCR or any other amplification technique prior to analysis. RNA, particularly mRNA, cDNA and genomic DNA may also be used in the same ways. Using amplification, characterization of the species and strain of infectious or resident organism present in an individual, may be made by an analysis of the genotype of a selected polynucleotide of the organism. Deletions and insertions can be detected by a change in size of the amplified product in comparison to a genotype of a reference sequence selected from a related organism, preferably a different species of the same genus or a different strain of the same species. Point mutations can be identified by hybridizing amplified DNA to labeled BASB027 polynucleotide sequences. Perfectly or significantly matched sequences can be distinguished from imperfectly or more significantly mismatched duplexes by DNase or RNase digestion, for DNA or RNA respectively, or by detecting differences in melting temperatures or renaturation kinetics. Polynucleotide sequence differences may also be detected by alterations in the electrophoretic mobility of polynucleotide fragments in gels as compared to a reference sequence. This may be carried out with or without denaturing agents. Polynucleotide differences may also be detected by direct DNA or RNA sequencing. See, for example, Myers *et al.*, *Science*, 230: 1242 (1985).

Sequence changes at specific locations also may be revealed by nuclease protection assays, such as RNase, V1 and S1 protection assay or a chemical cleavage method. See, for example, Cotton *et al.*, *Proc. Natl. Acad. Sci., USA*, 85: 4397-4401 (1985).

In another embodiment, an array of oligonucleotides probes comprising BASB027 nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of, for example, genetic mutations, serotype, taxonomic classification or identification. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability (see, for example, Chee *et al.*, *Science*, 274: 610 (1996)).

Thus in another aspect, the present invention relates to a diagnostic kit which comprises:

- (a) a polynucleotide of the present invention, preferably the nucleotide sequence of SEQ ID NO:1 or 3, or a fragment thereof;
- (b) a nucleotide sequence complementary to that of (a);
- (c) a polypeptide of the present invention, preferably the polypeptide of SEQ ID NO:2 or 4 or a fragment thereof; or
- (d) an antibody to a polypeptide of the present invention, preferably to the polypeptide of SEQ ID NO:2 or 4.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component. Such a kit will be of use in diagnosing a disease or susceptibility to a Disease, among others.

This invention also relates to the use of polynucleotides of the present invention as diagnostic reagents. Detection of a mutated form of a polynucleotide of the invention, preferably SEQ ID NO:1 or 3, which is associated with a disease or pathogenicity will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, a prognosis of a course of disease, a determination of a stage of disease, or a susceptibility to a disease.

which results from under-expression, over-expression or altered expression of the polynucleotide. Organisms, particularly infectious organisms, carrying mutations in such polynucleotide may be detected at the polynucleotide level by a variety of techniques, such as those described elsewhere herein.

Cells from an organism carrying mutations or polymorphisms (allelic variations) in a polynucleotide and/or polypeptide of the invention may also be detected at the polynucleotide or polypeptide level by a variety of techniques, to allow for serotyping, for example. For example, RT-PCR can be used to detect mutations in the RNA. It is particularly preferred to use RT-PCR in conjunction with automated detection systems, such as, for example, GeneScan. RNA, cDNA or genomic DNA may also be used for the same purpose. PCR. As an example, PCR primers complementary to a polynucleotide encoding BASB027 polypeptide can be used to identify and analyze mutations.

The invention further provides primers with 1, 2, 3 or 4 nucleotides removed from the 5' and/or the 3' end. These primers may be used for, among other things, amplifying BASB027 DNA and/or RNA isolated from a sample derived from an individual, such as a bodily material. The primers may be used to amplify a polynucleotide isolated from an infected individual, such that the polynucleotide may then be subject to various techniques for elucidation of the polynucleotide sequence. In this way, mutations in the polynucleotide sequence may be detected and used to diagnose and/or prognose the infection or its stage or course, or to serotype and/or classify the infectious agent.

The invention further provides a process for diagnosing, disease, preferably bacterial infections, more preferably infections caused by *Moraxella catarrhalis*, comprising determining from a sample derived from an individual, such as a bodily material, an increased level of expression of polynucleotide having a sequence of SEQ ID NO:1 or 3. Increased or decreased expression of a BASB027 polynucleotide can be measured using any one of the methods well known in the art for the quantitation of polynucleotides, such

as, for example, amplification, PCR, RT-PCR, RNase protection, Northern blotting, spectrometry and other hybridization methods.

In addition, a diagnostic assay in accordance with the invention for detecting over-expression of BASB027 polypeptide compared to normal control tissue samples may be used to detect the presence of an infection, for example. Assay techniques that can be used to determine levels of a BASB027 polypeptide, in a sample derived from a host, such as a bodily material, are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis, antibody sandwich assays, antibody detection and ELISA assays.

The polynucleotides of the invention may be used as components of polynucleotide arrays, preferably high density arrays or grids. These high density arrays are particularly useful for diagnostic and prognostic purposes. For example, a set of spots each comprising a different gene, and further comprising a polynucleotide or polynucleotides of the invention, may be used for probing, such as using hybridization or nucleic acid amplification, using a probes obtained or derived from a bodily sample, to determine the presence of a particular polynucleotide sequence or related sequence in an individual. Such a presence may indicate the presence of a pathogen, particularly *Moraxella catarrhalis*, and may be useful in diagnosing and/or prognosing disease or a course of disease. A grid comprising a number of variants of the polynucleotide sequence of SEQ ID NO:1 or 3 are preferred. Also preferred is a comprising a number of variants of a polynucleotide sequence encoding the polypeptide sequence of SEQ ID NO:2 or 4.

Antibodies

The polypeptides and polynucleotides of the invention or variants thereof, or cells expressing the same can be used as immunogens to produce antibodies immunospecific for such polypeptides or polynucleotides respectively.

In certain preferred embodiments of the invention there are provided antibodies against BASB027 polypeptides or polynucleotides.

Antibodies generated against the polypeptides or polynucleotides of the invention can be obtained by administering the polypeptides and/or polynucleotides of the invention, or epitope-bearing fragments of either or both, analogues of either or both, or cells expressing either or both, to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique known in the art that provides antibodies produced by continuous cell line cultures can be used. Examples include various techniques, such as those in Kohler, G. and Milstein, C., *Nature* 256: 495-497 (1975); Kozbor *et al.*, *Immunology Today* 4: 72 (1983); Cole *et al.*, pg. 77-96 in *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc. (1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can be adapted to produce single chain antibodies to polypeptides or polynucleotides of this invention. Also, transgenic mice, or other organisms or animals, such as other mammals, may be used to express humanized antibodies immunospecific to the polypeptides or polynucleotides of the invention.

Alternatively, phage display technology may be utilized to select antibody genes with binding activities towards a polypeptide of the invention either from repertoires of PCR amplified v-genes of lymphocytes from humans screened for possessing anti-BASB027 or from naive libraries (McCafferty, *et al.*, (1990), *Nature* 348, 552-554; Marks, *et al.*, (1992) *Biotechnology* 10, 779-783). The affinity of these antibodies can also be improved by, for example, chain shuffling (Clackson *et al.*, (1991) *Nature* 352: 628).

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptides or polynucleotides of the invention to purify the polypeptides or polynucleotides by, for example, affinity chromatography.

Thus, among others, antibodies against BASB027-polypeptide or BASB027-polynucleotide may be employed to treat infections, particularly bacterial infections.

Polypeptide variants include antigenically, epitopically or immunologically equivalent variants form a particular aspect of this invention.

Preferably, the antibody or variant thereof is modified to make it less immunogenic in the individual. For example, if the individual is human the antibody may most preferably be "humanized," where the complementarity determining region or regions of the hybridoma-derived antibody has been transplanted into a human monoclonal antibody, for example as described in Jones *et al.* (1986), *Nature* 321, 522-525 or Tempest *et al.*, (1991) *Biotechnology* 9, 266-273.

Antagonists and Agonists - Assays and Molecules

Polypeptides and polynucleotides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See, *e.g.*, Coligan *et al.*, *Current Protocols in Immunology* 1(2): Chapter 5 (1991).

The screening methods may simply measure the binding of a candidate compound to the polypeptide or polynucleotide, or to cells or membranes bearing the polypeptide or polynucleotide, or a fusion protein of the polypeptide by means of a label directly or indirectly associated with the candidate compound. Alternatively, the screening method

may involve competition with a labeled competitor. Further, these screening methods may test whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide or polynucleotide, using detection systems appropriate to the cells comprising the polypeptide or polynucleotide. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed. Constitutively active polypeptide and/or constitutively expressed polypeptides and polynucleotides may be employed in screening methods for inverse agonists or inhibitors, in the absence of an agonist or inhibitor, by testing whether the candidate compound results in inhibition of activation of the polypeptide or polynucleotide, as the case may be. Further, the screening methods may simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide or polynucleotide of the present invention, to form a mixture, measuring BASB027 polypeptide and/or polynucleotide activity in the mixture, and comparing the BASB027 polypeptide and/or polynucleotide activity of the mixture to a standard. Fusion proteins, such as those made from Fc portion and BASB027 polypeptide, as hereinbefore described, can also be used for high-throughput screening assays to identify antagonists of the polypeptide of the present invention, as well as of phylogenetically and and/or functionally related polypeptides (see D. Bennett *et al.*, J Mol Recognition, 8:52-58 (1995); and K. Johanson *et al.*, J Biol Chem, 270(16):9459-9471 (1995)).

The polynucleotides, polypeptides and antibodies that bind to and/or interact with a polypeptide of the present invention may also be used to configure screening methods for detecting the effect of added compounds on the production of mRNA and/or polypeptide in cells. For example, an ELISA assay may be constructed for measuring secreted or cell associated levels of polypeptide using monoclonal and polyclonal antibodies by standard methods known in the art. This can be used to discover agents which may inhibit or enhance the production of polypeptide (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues.

The invention also provides a method of screening compounds to identify those which enhance (agonist) or block (antagonist) the action of BASB027 polypeptides or polynucleotides, particularly those compounds that are bacteriostatic and/or bactericidal. The method of screening may involve high-throughput techniques. For example, to screen for agonists or antagonists, a synthetic reaction mix, a cellular compartment, such as a membrane, cell envelope or cell wall, or a preparation of any thereof, comprising BASB027 polypeptide and a labeled substrate or ligand of such polypeptide is incubated in the absence or the presence of a candidate molecule that may be a BASB027 agonist or antagonist. The ability of the candidate molecule to agonize or antagonize the BASB027 polypeptide is reflected in decreased binding of the labeled ligand or decreased production of product from such substrate. Molecules that bind gratuitously, *i.e.*, without inducing the effects of BASB027 polypeptide are most likely to be good antagonists. Molecules that bind well and, as the case may be, increase the rate of product production from substrate, increase signal transduction, or increase chemical channel activity are agonists. Detection of the rate or level of, as the case may be, production of product from substrate, signal transduction, or chemical channel activity may be enhanced by using a reporter system. Reporter systems that may be useful in this regard include but are not limited to colorimetric, labeled substrate converted into product, a reporter gene that is responsive to changes in BASB027 polynucleotide or polypeptide activity, and binding assays known in the art.

Another example of an assay for BASB027 agonists is a competitive assay that combines BASB027 and a potential agonist with BASB027-binding molecules, recombinant BASB027 binding molecules, natural substrates or ligands, or substrate or ligand mimetics, under appropriate conditions for a competitive inhibition assay. BASB027 can be labeled, such as by radioactivity or a colorimetric compound, such that the number of BASB027 molecules bound to a binding molecule or converted to product can be determined accurately to assess the effectiveness of the potential antagonist.

Potential antagonists include, among others, small organic molecules, peptides, polypeptides and antibodies that bind to a polynucleotide and/or polypeptide of the invention and thereby inhibit or extinguish its activity or expression. Potential antagonists also may be small organic molecules, a peptide, a polypeptide such as a closely related protein or antibody that binds the same sites on a binding molecule, such as a binding molecule, without inducing BASB027-induced activities, thereby preventing the action or expression of BASB027 polypeptides and/or polynucleotides by excluding BASB027 polypeptides and/or polynucleotides from binding.

Potential antagonists include a small molecule that binds to and occupies the binding site of the polypeptide thereby preventing binding to cellular binding molecules, such that normal biological activity is prevented. Examples of small molecules include but are not limited to small organic molecules, peptides or peptide-like molecules. Other potential antagonists include antisense molecules (see Okano, *J. Neurochem.* 56: 560 (1991); *OLIGODEOXYNUCLEOTIDES AS ANTISENSE INHIBITORS OF GENE EXPRESSION*, CRC Press, Boca Raton, FL (1988), for a description of these molecules). Preferred potential antagonists include compounds related to and variants of BASB027.

In a further aspect, the present invention relates to genetically engineered soluble fusion proteins comprising a polypeptide of the present invention, or a fragment thereof, and various portions of the constant regions of heavy or light chains of immunoglobulins of various subclasses (IgG, IgM, IgA, IgE). Preferred as an immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG1, where fusion takes place at the hinge region. In a particular embodiment, the Fc part can be removed simply by incorporation of a cleavage sequence which can be cleaved with blood clotting factor Xa. Furthermore, this invention relates to processes for the preparation of these fusion proteins by genetic engineering, and to the use thereof for drug screening, diagnosis and therapy. A further aspect of the invention also relates to polynucleotides encoding such

fusion proteins. Examples of fusion protein technology can be found in International Patent Application Nos. WO94/29458 and WO94/22914.

Each of the polynucleotide sequences provided herein may be used in the discovery and development of antibacterial compounds. The encoded protein, upon expression, can be used as a target for the screening of antibacterial drugs. Additionally, the polynucleotide sequences encoding the amino terminal regions of the encoded protein or Shine-Delgarno or other translation facilitating sequences of the respective mRNA can be used to construct antisense sequences to control the expression of the coding sequence of interest.

The invention also provides the use of the polypeptide, polynucleotide, agonist or antagonist of the invention to interfere with the initial physical interaction between a pathogen or pathogens and a eukaryotic, preferably mammalian, host responsible for sequelae of infection. In particular, the molecules of the invention may be used: in the prevention of adhesion of bacteria, in particular gram positive and/or gram negative bacteria, to eukaryotic, preferably mammalian, extracellular matrix proteins on in-dwelling devices or to extracellular matrix proteins in wounds; to block bacterial adhesion between eukaryotic, preferably mammalian, extracellular matrix proteins and bacterial BASB027 proteins that mediate tissue damage and/or; to block the normal progression of pathogenesis in infections initiated other than by the implantation of in-dwelling devices or by other surgical techniques.

In accordance with yet another aspect of the invention, there are provided BASB027 agonists and antagonists, preferably bacteristatic or bactericidal agonists and antagonists.

The antagonists and agonists of the invention may be employed, for instance, to prevent, inhibit and/or treat diseases.

In a further aspect, the present invention relates to mimotopes of the polypeptide of the invention. A mimotope is a peptide sequence, sufficiently similar to the native peptide (sequentially or structurally), which is capable of being recognised by antibodies which recognise the native peptide; or is capable of raising antibodies which recognise the native peptide when coupled to a suitable carrier.

Peptide mimotopes may be designed for a particular purpose by addition, deletion or substitution of elected amino acids. Thus, the peptides may be modified for the purposes of ease of conjugation to a protein carrier. For example, it may be desirable for some chemical conjugation methods to include a terminal cysteine. In addition it may be desirable for peptides conjugated to a protein carrier to include a hydrophobic terminus distal from the conjugated terminus of the peptide, such that the free unconjugated end of the peptide remains associated with the surface of the carrier protein. Thereby presenting the peptide in a conformation which most closely resembles that of the peptide as found in the context of the whole native molecule. For example, the peptides may be altered to have an N-terminal cysteine and a C-terminal hydrophobic amidated tail. Alternatively, the addition or substitution of a D-stereoisomer form of one or more of the amino acids may be performed to create a beneficial derivative, for example to enhance stability of the peptide.

Alternatively, peptide mimotopes may be identified using antibodies which are capable themselves of binding to the polypeptides of the present invention using techniques such as phage display technology (EP 0 552 267 B1). This technique, generates a large number of peptide sequences which mimic the structure of the native peptides and are, therefore, capable of binding to anti-native peptide antibodies, but may not necessarily themselves share significant sequence homology to the native polypeptide.

Vaccines

Another aspect of the invention relates to a method for inducing an immunological response in an individual, particularly a mammal, preferably humans, which comprises inoculating the individual with BASB027 polynucleotide and/or polypeptide, or a fragment or variant thereof, adequate to produce antibody and/ or T cell immune response to protect said individual from infection, particularly bacterial infection and most particularly *Moraxella catarrhalis* infection. Also provided are methods whereby such immunological response slows bacterial replication. Yet another aspect of the invention relates to a method of inducing immunological response in an individual which comprises delivering to such individual a nucleic acid vector, sequence or ribozyme to direct expression of BASB027 polynucleotide and/or polypeptide, or a fragment or a variant thereof, for expressing BASB027 polynucleotide and/or polypeptide, or a fragment or a variant thereof *in vivo* in order to induce an immunological response, such as, to produce antibody and/ or T cell immune response, including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said individual, preferably a human, from disease, whether that disease is already established within the individual or not. One example of administering the gene is by accelerating it into the desired cells as a coating on particles or otherwise. Such nucleic acid vector may comprise DNA, RNA, a ribozyme, a modified nucleic acid, a DNA/RNA hybrid, a DNA-protein complex or an RNA-protein complex.

A further aspect of the invention relates to an immunological composition that when introduced into an individual, preferably a human, capable of having induced within it an immunological response, induces an immunological response in such individual to a BASB027 polynucleotide and/or polypeptide encoded therefrom, wherein the composition comprises a recombinant BASB027 polynucleotide and/or polypeptide encoded therefrom and/or comprises DNA and/or RNA which encodes and expresses an antigen of said BASB027 polynucleotide, polypeptide encoded therefrom, or other polypeptide of the invention. The immunological response may be used therapeutically or prophylactically and may take the form of antibody immunity and/or cellular immunity, such as cellular immunity arising from CTL or CD4+ T cells.

A BASB027 polypeptide or a fragment thereof may be fused with co-protein or chemical moiety which may or may not by itself produce antibodies, but which is capable of stabilizing the first protein and producing a fused or modified protein which will have antigenic and/or immunogenic properties, and preferably protective properties. Thus fused recombinant protein, preferably further comprises an antigenic co-protein, such as lipoprotein D from *Haemophilus influenzae*, Glutathione-S-transferase (GST) or beta-galactosidase, or any other relatively large co-protein which solubilizes the protein and facilitates production and purification thereof. Moreover, the co-protein may act as an adjuvant in the sense of providing a generalized stimulation of the immune system of the organism receiving the protein. The co-protein may be attached to either the amino- or carboxy-terminus of the first protein.

Provided by this invention are compositions, particularly vaccine compositions, and methods comprising the polypeptides and/or polynucleotides of the invention and immunostimulatory DNA sequences, such as those described in Sato, Y. *et al.* Science 273: 352 (1996).

Also, provided by this invention are methods using the described polynucleotide or particular fragments thereof, which have been shown to encode non-variable regions of bacterial cell surface proteins, in polynucleotide constructs used in such genetic immunization experiments in animal models of infection with *Moraxella catarrhalis*. Such experiments will be particularly useful for identifying protein epitopes able to provoke a prophylactic or therapeutic immune response. It is believed that this approach will allow for the subsequent preparation of monoclonal antibodies of particular value, derived from the requisite organ of the animal successfully resisting or clearing infection, for the development of prophylactic agents or therapeutic treatments of bacterial infection, particularly *Moraxella catarrhalis* infection, in mammals, particularly humans.

The invention also includes a vaccine formulation which comprises an immunogenic recombinant polypeptide and/or polynucleotide of the invention together with a suitable carrier, such as a pharmaceutically acceptable carrier. Since the polypeptides and polynucleotides may be broken down in the stomach, each is preferably administered parenterally, including, for example, administration that is subcutaneous, intramuscular, intravenous, or intradermal. Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostatic compounds and solutes which render the formulation isotonic with the bodily fluid, preferably the blood, of the individual; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use.

The vaccine formulation of the invention may also include adjuvant systems for enhancing the immunogenicity of the formulation. Preferably the adjuvant system raises preferentially a TH1 type of response.

An immune response may be broadly distinguished into two extreme categories, being a humoral or cell mediated immune responses (traditionally characterised by antibody and cellular effector mechanisms of protection respectively). These categories of response have been termed TH1-type responses (cell-mediated response), and TH2-type immune responses (humoral response).

Extreme TH1-type immune responses may be characterised by the generation of antigen specific, haplotype restricted cytotoxic T lymphocytes, and natural killer cell responses. In mice TH1-type responses are often characterised by the generation of antibodies of the IgG2a subtype, whilst in the human these correspond to IgG1 type antibodies. TH2-

type immune responses are characterised by the generation of a broad range of immunoglobulin isotypes including in mice IgG1, IgA, and IgM.

It can be considered that the driving force behind the development of these two types of immune responses are cytokines. High levels of TH1-type cytokines tend to favour the induction of cell mediated immune responses to the given antigen, whilst high levels of TH2-type cytokines tend to favour the induction of humoral immune responses to the antigen.

The distinction of TH1 and TH2-type immune responses is not absolute. In reality an individual will support an immune response which is described as being predominantly TH1 or predominantly TH2. However, it is often convenient to consider the families of cytokines in terms of that described in murine CD4 +ve T cell clones by Mosmann and Coffman (Mosmann, T.R. and Coffman, R.L. (1989) *TH1 and TH2 cells: different patterns of lymphokine secretion lead to different functional properties. Annual Review of Immunology*, 7, p145-173). Traditionally, TH1-type responses are associated with the production of the INF- γ and IL-2 cytokines by T-lymphocytes. Other cytokines often directly associated with the induction of TH1-type immune responses are not produced by T-cells, such as IL-12. In contrast, TH2- type responses are associated with the secretion of IL-4, IL-5, IL-6 and IL-13.

It is known that certain vaccine adjuvants are particularly suited to the stimulation of either TH1 or TH2 - type cytokine responses. Traditionally the best indicators of the TH1:TH2 balance of the immune response after a vaccination or infection includes direct measurement of the production of TH1 or TH2 cytokines by T lymphocytes *in vitro* after restimulation with antigen, and/or the measurement of the IgG1:IgG2a ratio of antigen specific antibody responses.

Thus, a TH1-type adjuvant is one which preferentially stimulates isolated T-cell populations to produce high levels of TH1-type cytokines when re-stimulated with antigen *in vitro*, and promotes development of both CD8+ cytotoxic T lymphocytes and antigen specific immunoglobulin responses associated with TH1-type isotype.

Adjuvants which are capable of preferential stimulation of the TH1 cell response are described in International Patent Application No. WO 94/00153 and WO 95/17209.

3 De-O-acylated monophosphoryl lipid A (3D-MPL) is one such adjuvant. This is known from GB 2220211 (Ribi). Chemically it is a mixture of 3 De-O-acylated monophosphoryl lipid A with 4, 5 or 6 acylated chains and is manufactured by Ribi Immunochem, Montana. A preferred form of 3 De-O-acylated monophosphoryl lipid A is disclosed in European Patent 0 689 454 B1 (SmithKline Beecham Biologicals SA).

Preferably, the particles of 3D-MPL are small enough to be sterile filtered through a 0.22micron membrane (European Patent number 0 689 454).

3D-MPL will be present in the range of 10µg - 100µg preferably 25-50µg per dose wherein the antigen will typically be present in a range 2-50µg per dose.

Another preferred adjuvant comprises QS21, an Hplc purified non-toxic fraction derived from the bark of *Quillaja Saponaria Molina*. Optionally this may be admixed with 3 De-O-acylated monophosphoryl lipid A (3D-MPL), optionally together with an carrier.

The method of production of QS21 is disclosed in US patent No. 5,057,540.

Non-reactogenic adjuvant formulations containing QS21 have been described previously (WO 96/33739). Such formulations comprising QS21 and cholesterol have been shown to be successful TH1 stimulating adjuvants when formulated together with an antigen.

Further adjuvants which are preferential stimulators of TH1 cell response include immunomodulatory oligonucleotides, for example unmethylated CpG sequences as disclosed in WO 96/02555.

Combinations of different TH1 stimulating adjuvants, such as those mentioned hereinabove, are also contemplated as providing an adjuvant which is a preferential stimulator of TH1 cell response. For example, QS21 can be formulated together with 3D-MPL. The ratio of QS21 : 3D-MPL will typically be in the order of 1 : 10 to 10 : 1; preferably 1:5 to 5 : 1 and often substantially 1 : 1. The preferred range for optimal synergy is 2.5 : 1 to 1 : 1 3D-MPL: QS21.

Preferably a carrier is also present in the vaccine composition according to the invention. The carrier may be an oil in water emulsion, or an aluminium salt, such as aluminium phosphate or aluminium hydroxide.

A preferred oil-in-water emulsion comprises a metabolisable oil, such as squalene, alpha tocopherol and Tween 80. In a particularly preferred aspect the antigens in the vaccine composition according to the invention are combined with QS21 and 3D-MPL in such an emulsion. Additionally the oil in water emulsion may contain span 85 and/or lecithin and/or tricaprylin.

Typically for human administration QS21 and 3D-MPL will be present in a vaccine in the range of 1µg - 200µg, such as 10-100µg, preferably 10µg - 50µg per dose.

Typically the oil in water will comprise from 2 to 10% squalene, from 2 to 10% alpha tocopherol and from 0.3 to 3% tween 80. Preferably the ratio of squalene: alpha tocopherol is equal to or less than 1 as this provides a more stable emulsion. Span 85 may also be present at a level of 1%. In some cases it may be advantageous that the vaccines of the present invention will further contain a stabiliser.

Non-toxic oil in water emulsions preferably contain a non-toxic oil, e.g. squalane or squalene, an emulsifier, e.g. Tween 80, in an aqueous carrier. The aqueous carrier may be, for example, phosphate buffered saline.

A particularly potent adjuvant formulation involving QS21, 3D-MPL and tocopherol in an oil in water emulsion is described in WO 95/17210.

The present invention also provides a polyvalent vaccine composition comprising a vaccine formulation of the invention in combination with other antigens, in particular antigens useful for treating cancers, autoimmune diseases and related conditions. Such a polyvalent vaccine composition may include a TH-1 inducing adjuvant as hereinbefore described.

While the invention has been described with reference to certain BASB027 polypeptides and polynucleotides, it is to be understood that this covers fragments of the naturally occurring polypeptides and polynucleotides, and similar polypeptides and polynucleotides with additions, deletions or substitutions which do not substantially affect the immunogenic properties of the recombinant polypeptides or polynucleotides.

Compositions, kits and administration

In a further aspect of the invention there are provided compositions comprising a BASB027 polynucleotide and/or a BASB027 polypeptide for administration to a cell or to a multicellular organism.

The invention also relates to compositions comprising a polynucleotide and/or a polypeptides discussed herein or their agonists or antagonists. The polypeptides and polynucleotides of the invention may be employed in combination with a non-sterile or sterile carrier or carriers for use with cells, tissues or organisms, such as a pharmaceutical

carrier suitable for administration to an individual. Such compositions comprise, for instance, a media additive or a therapeutically effective amount of a polypeptide and/or polynucleotide of the invention and a pharmaceutically acceptable carrier or excipient. Such carriers may include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol and combinations thereof. The formulation should suit the mode of administration. The invention further relates to diagnostic and pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

Polypeptides, polynucleotides and other compounds of the invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

The pharmaceutical compositions may be administered in any effective, convenient manner including, for instance, administration by topical, oral, anal, vaginal, intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal or intradermal routes among others.

In therapy or as a prophylactic, the active agent may be administered to an individual as an injectable composition, for example as a sterile aqueous dispersion, preferably isotonic.

In a further aspect, the present invention provides for pharmaceutical compositions comprising a therapeutically effective amount of a polypeptide and/or polynucleotide, such as the soluble form of a polypeptide and/or polynucleotide of the present invention, agonist or antagonist peptide or small molecule compound, in combination with a pharmaceutically acceptable carrier or excipient. Such carriers include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

Polypeptides, polynucleotides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

The composition will be adapted to the route of administration, for instance by a systemic or an oral route. Preferred forms of systemic administration include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if a polypeptide or other compounds of the present invention can be formulated in an enteric or an encapsulated formulation, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels, solutions, powders and the like.

For administration to mammals, and particularly humans, it is expected that the daily dosage level of the active agent will be from 0.01 mg/kg to 10 mg/kg, typically around 1 mg/kg. The physician in any event will determine the actual dosage which will be most suitable for an individual and will vary with the age, weight and response of the particular individual. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

The dosage range required depends on the choice of peptide, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages, however, are in the range of 0.1-100 μ g/kg of subject.

A vaccine composition is conveniently in injectable form. Conventional adjuvants may be employed to enhance the immune response. A suitable unit dose for vaccination is 0.5-5 microgram/kg of antigen, and such dose is preferably administered 1-3 times and with an

interval of 1-3 weeks. With the indicated dose range, no adverse toxicological effects will be observed with the compounds of the invention which would preclude their administration to suitable individuals.

Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

Sequence Databases, Sequences in a Tangible Medium, and Algorithms

Polynucleotide and polypeptide sequences form a valuable information resource with which to determine their 2- and 3-dimensional structures as well as to identify further sequences of similar homology. These approaches are most easily facilitated by storing the sequence in a computer readable medium and then using the stored data in a known macromolecular structure program or to search a sequence database using well known searching tools, such as the GCG program package.

Also provided by the invention are methods for the analysis of character sequences or strings, particularly genetic sequences or encoded protein sequences. Preferred methods of sequence analysis include, for example, methods of sequence homology analysis, such as identity and similarity analysis, DNA, RNA and protein structure analysis, sequence assembly, cladistic analysis, sequence motif analysis, open reading frame determination, nucleic acid base calling, codon usage analysis, nucleic acid base trimming, and sequencing chromatogram peak analysis.

A computer based method is provided for performing homology identification. This method comprises the steps of: providing a first polynucleotide sequence comprising the

sequence of a polynucleotide of the invention in a computer readable medium; and comparing said first polynucleotide sequence to at least one second polynucleotide or polypeptide sequence to identify homology.

A computer based method is also provided for performing homology identification. said method comprising the steps of: providing a first polypeptide sequence comprising the sequence of a polypeptide of the invention in a computer readable medium; and comparing said first polypeptide sequence to at least one second polynucleotide or polypeptide sequence to identify homology.

All publications and references, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference in their entirety as if each individual publication or reference were specifically and individually indicated to be incorporated by reference herein as being fully set forth. Any patent application to which this application claims priority is also incorporated by reference herein in its entirety in the manner described above for publications and references.

DEFINITIONS

"Identity," as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as the case may be, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" can be readily calculated by known methods, including but not limited to those described in (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data*, Part I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heine,

G., Academic Press. 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press. New York. 1991; and Carillo. H., and Lipman, D., *SIAM J. Applied Math.*, 48: 1073 (1988). Methods to determine identity are designed to give the largest match between the sequences tested. Moreover, methods to determine identity are codified in publicly available computer programs. Computer program methods to determine identity between two sequences include, but are not limited to, the GAP program in the GCG program package (Devereux, J., et al., *Nucleic Acids Research* 12(1): 387 (1984)), BLASTP, BLASTN (Altschul, S.F. et al., *J. Molec. Biol.* 215: 403-410 (1990)), and FASTA (Pearson and Lipman *Proc. Natl. Acad. Sci. USA* 85: 2444-2448 (1988)). The BLAST family of programs is publicly available from NCBI and other sources (*BLAST Manual*, Altschul, S., et al., NCBI NLM NIH Bethesda, MD 20894; Altschul, S., et al., *J. Mol. Biol.* 215: 403-410 (1990)). The well known Smith Waterman algorithm may also be used to determine identity.

Parameters for polypeptide sequence comparison include the following:

Algorithm: Needleman and Wunsch, *J. Mol Biol.* 48: 443-453 (1970)

Comparison matrix: BLOSSUM62 from Henikoff and Henikoff,
Proc. Natl. Acad. Sci. USA. 89:10915-10919 (1992)

Gap Penalty: 8

Gap Length Penalty: 2

A program useful with these parameters is publicly available as the "gap" program from Genetics Computer Group, Madison WI. The aforementioned parameters are the default parameters for peptide comparisons (along with no penalty for end gaps).

Parameters for polynucleotide comparison include the following:

Algorithm: Needleman and Wunsch, *J. Mol Biol.* 48: 443-453 (1970)

Comparison matrix: matches = +10, mismatch = 0

Gap Penalty: 50

Gap Length Penalty: 3

Available as: The "gap" program from Genetics Computer Group, Madison WI. These are the default parameters for nucleic acid comparisons.

A preferred meaning for "identity" for polynucleotides and polypeptides, as the case may be, are provided in (1) and (2) below.

(1) Polynucleotide embodiments further include an isolated polynucleotide comprising a polynucleotide sequence having at least a 50, 60, 70, 80, 85, 90, 95, 97 or 100% identity to the reference sequence of SEQ ID NO:1, wherein said polynucleotide sequence may be identical to the reference sequence of SEQ ID NO:1 or may include up to a certain integer number of nucleotide alterations as compared to the reference sequence, wherein said alterations are selected from the group consisting of at least one nucleotide deletion, substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence, and wherein said number of nucleotide alterations is determined by multiplying the total number of nucleotides in SEQ ID NO:1 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of nucleotides in SEQ ID NO:1, or:

$$n_n \leq x_n - (x_n \bullet y),$$

wherein n_n is the number of nucleotide alterations, x_n is the total number of nucleotides in SEQ ID NO:1, y is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and \bullet is the symbol for the multiplication operator, and wherein any non-integer product of x_n and y is rounded down to the nearest integer prior to subtracting it from x_n . Alterations of a polynucleotide sequence encoding the polypeptide of SEQ ID NO:2 may create nonsense, missense or

frameshift mutations in this coding sequence and thereby alter the polypeptide encoded by the polynucleotide following such alterations.

By way of example, a polynucleotide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:1, that is it may be 100% identical, or it may include up to a certain integer number of nucleic acid alterations as compared to the reference sequence such that the percent identity is less than 100% identity. Such alterations are selected from the group consisting of at least one nucleic acid deletion, substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal positions of the reference polynucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleic acids in the reference sequence or in one or more contiguous groups within the reference sequence. The number of nucleic acid alterations for a given percent identity is determined by multiplying the total number of nucleic acids in SEQ ID NO:1 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of nucleic acids in SEQ ID NO:1, or:

$$n_n \leq x_n - (x_n \bullet y),$$

wherein n_n is the number of nucleic acid alterations, x_n is the total number of nucleic acids in SEQ ID NO:1, y is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., \bullet is the symbol for the multiplication operator, and wherein any non-integer product of x_n and y is rounded down to the nearest integer prior to subtracting it from x_n .

(2) Polypeptide embodiments further include an isolated polypeptide comprising a polypeptide having at least a 50, 60, 70, 80, 85, 90, 95, 97 or 100% identity to a polypeptide reference sequence of SEQ ID NO:2, wherein said polypeptide sequence may be identical to the reference sequence of SEQ ID NO:2 or may include up to a certain integer number of amino acid alterations as compared to the reference sequence, wherein

said alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence, and wherein said number of amino acid alterations is determined by multiplying the total number of amino acids in SEQ ID NO:2 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

$$n_a \leq x_a - (x_a \bullet y),$$

wherein n_a is the number of amino acid alterations, x_a is the total number of amino acids in SEQ ID NO:2, y is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and \bullet is the symbol for the multiplication operator, and wherein any non-integer product of x_a and y is rounded down to the nearest integer prior to subtracting it from x_a .

By way of example, a polypeptide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:2, that is it may be 100% identical, or it may include up to a certain integer number of amino acid alterations as compared to the reference sequence such that the percent identity is less than 100% identity. Such alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence. The number of amino acid alterations for a given % identity is determined by multiplying the total number of amino

acids in SEQ ID NO:2 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

$$n_a \leq x_a - (x_a \bullet y),$$

wherein n_a is the number of amino acid alterations, x_a is the total number of amino acids in SEQ ID NO:2, y is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., and \bullet is the symbol for the multiplication operator, and wherein any non-integer product of x_a and y is rounded down to the nearest integer prior to subtracting it from x_a .

"Individual(s)," when used herein with reference to an organism, means a multicellular eukaryote, including, but not limited to a metazoan, a mammal, an ovid, a bovid, a simian, a primate, and a human.

"Isolated" means altered "by the hand of man" from its natural state, *i.e.*, if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein. Moreover, a polynucleotide or polypeptide that is introduced into an organism by transformation, genetic manipulation or by any other recombinant method is "isolated" even if it is still present in said organism, which organism may be living or non-living.

"Polynucleotide(s)" generally refers to any polyribonucleotide or polydeoxyribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA including single and double-stranded regions.

"Variant" refers to a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide, but retains essential properties. A typical variant of a

polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

"Disease(s)" means any disease caused by or related to infection by a bacteria, including, for example, otitis media in infants and children, pneumonia in elderlies, sinusitis, nosocomial infections and invasive diseases, chronic otitis media with hearing loss, fluid accumulation in the middle ear, auditive nerve damage, delayed speech learning, infection of the upper respiratory tract and inflammation of the middle ear.

EXAMPLES:

The examples below are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The examples are illustrative, but do not limit the invention.

Example 1:**Discovery and confirmatory DNA sequencing of the BASB027 gene from *Moraxella catarrhalis* strain ATCC 43617.**

The BASB027 gene of SEQ ID NO:1 was first discovered in the Incyte PathoSeq data base containing unfinished genomic DNA sequences of the *Moraxella catarrhalis* strain ATCC 43617 (also referred to as strain Mc2931). The translation of the BASB027 polynucleotide sequence, shown in SEQ ID NO:2, showed significant similarity (32 % identity in a 817 amino acids overlap) to the OMP85 outer membrane protein of *Neisseria meningitidis*.

The sequence of the BASB027 gene was further confirmed experimentally. For this purpose, genomic DNA was extracted from 10^{10} cells of the *M. catarrhalis* cells (strain ATCC 43617) using the QIAGEN genomic DNA extraction kit (Qiagen GmbH), and 1 µg of this material was submitted to Polymerase Chain Reaction DNA amplification using primers E515515 (5'- ACT-ATA-GGG-CAC-GCG-TG -3') [SEQ ID NO:5] and E515528 : (5'- CCT-GCG-TTT-GTT-TGA-TTG-AG-3') [SEQ ID NO:6]. This PCR product was purified on a Biorobot 9600 (Qiagen GmbH) apparatus and subjected to DNA sequencing using the Big Dye Cycle Sequencing kit (Perkin-Elmer) and an ABI 377/PRISM DNA sequencer. DNA sequencing was performed on both strands with a redundancy of 2 and the full length sequence was assembled using the SeqMan program from the DNASTAR Lasergene software package. The resulting DNA sequence and deduced polypeptide sequence are shown as SEQ ID NO:3 and SEQ ID NO:4

respectively. Four nucleotide differences distinguish SEQ ID NO:3 from SEQ ID NO:1. Using the MEGALIGN program from the DNASTAR Lasergene software package, an alignment of the polynucleotide sequences of SEQ ID NO:1 and 3 was performed, and is displayed in Figure 2; their level of identity was calculated to be 99.8 %. Using the same program, an alignment of the polypeptide sequences of SEQ ID NO:2 and 4 was performed, and is displayed in Figure 3; their level of identity was calculated to be 99.8 %.

Example 2:

Variability analysis of the BASB027 gene among several *Moraxella catarrhalis* strains.

2A: Restriction Fragment Length Analysis (RFLP).

Genomic DNA was extracted from 16 *M. catarrhalis* strains (presented in Table 1) as described below. *M. catarrhalis* was streaked for single colonies on BHI agar plates and grown overnight at 37 °C. Three or four single colonies were picked and used to inoculate a ~1.5 ml BHI (Brain-heart infusion) broth seed culture which was grown overnight in a shaking incubator, ~300 rpm, at 37 °C. A 500ml erlenmeyer flask containing ~150 ml of BHI broth was inoculated with the seed culture and grown for ~12-16 hours at 37 °C in a shaking incubator, ~175 rpm, to generate cell mass for DNA isolation. Cells were collected by centrifugation in a Sorvall GSA rotor at ~2000 X g for 15 minutes at room temperature. The supernatant was removed and the cell pellet suspended in ~5.0 ml of sterile water. An equal volume of lysis buffer (200 mM NaCl, 20 mM EDTA, 40 mM Tris-Hcl, pH 8.0, 0.5% (w/v) SDS, 0.5% (v/v) 2-mercaptoethanol, and 250 µg/ml of proteinase K) was added and the cells suspended by gentle agitation and trituration. The cell suspension was then incubated ~12 hours at 50°C to lyse the bacteria and liberate chromosomal DNA. Proteinaceous material was precipitated by the addition of 5.0 ml of saturated NaCl (~6.0 M, in sterile water) and centrifugation at ~5,500xg in a Sorvall SS34 rotor at room temperature. Chromosomal

DNA was precipitated from the cleared supernatant by the addition of two volumes of 100 % ethanol. Aggregated DNA was collected and washed using gentle agitation in a small volume of a 70 % ethanol solution. Purified chromosomal DNA was suspended in sterile water and allowed to dissolve/ disburse overnight at 4 °C by gentle rocking. The concentration of dissolved DNA was determined spectrophotometrically at 260 nm using an extinction coefficient of 1.0 O.D. unit ~50 µg/ml.

This material was next submitted to PCR amplification using the MC-D15-BamF (5'-AAG GGC CCA ATT ACG CAG AGG GGA TCC ACA GGA CTA CAG CGA GTG ACC ATT GAA AGC TTA C -3') [SEQ ID NO:7] and MC-D15-SalRC (AAG GGC CCA ATT ACG CAG AGG GTC GAC TTA TTA AAA GAC ACT ACC AAT CTG GAA CTG TAC CGT ATC G -3') [SEQ ID NO:8] oligonucleotides. The corresponding BASB027 gene amplicons were then subjected independantly to hydrolysis using restriction enzymes (*AciI*, *HindIII*, *MaeIII*, *NlaIII*, *RsaI*, *Sau3AI*) and restriction products were separated by agarose or polyacrylamide gel electrophoresis using standard molecular biology procedures as described in "Molecular Cloning, a Laboratory Manual, Second Edition, Eds: Sambrook, Fritsch & Maniatis, Cold Spring Harbor press 1989". The photographs of the resulting electrophoresis gels are displayed in Figure 1. For each strain, RFLP patterns corresponding to the 6 restriction enzymes were scored and combined. Groups of strains sharing identical combination of RFLP patterns were then defined. Using this methodology, the strains tested in this study fell into 4 genomic groups (Group 1: Mc2906, Mc 2908, Mc2912, Mc2926; Group 2: Mc2905, Mc2907, Mc2909, Mc2911, Mc2913, Mc2960, Mc2975 ; Group 3: Mc2910, Mc2912, Mc2956, Mc2969; Group 4: Mc2931). These data support that the *Moraxella catarrhalis* population used in this study displays limited nucleotide sequence diversity for the BASB027 gene.

Table 1: Features of the *Moraxella catarrhalis* strains used in this study

Strain	Isolated in:	from:
Mc2904	USA	Tympanocentesis
Mc2905	USA	Tympanocentesis
Mc2906	USA	Tympanocentesis
Mc2907	USA	Tympanocentesis
Mc2908	USA	Acute otitis Tympanocentesis
Mc2909	USA	Tympanocentesis
Mc2910	USA	Tympanocentesis
Mc2911	USA	Acute otitis Tympanocentesis
Mc2912	USA	Acute otitis Tympanocentesis
Mc2913	USA	Acute otitis Tympanocentesis
Mc2926	USA	Tympanocentesis
Mc2931 /ATCC 43617	USA	Transtracheal aspirate
Mc2956	Finland	Middle ear fluid
Mc2960	Finland	Middle ear fluid
Mc2969	Norway	Nasopharynx (Pharyngitis- Rhinitis)
Mc2975	Norway	Nasopharynx (Rhinitis)

Example 3: Construction of Plasmid to Express Recombinant BASB027

A: Cloning of BASB027.

The *Bam*HI and *Sal*I restriction sites engineered into the forward ([SEQ ID NO:7]) and reverse complementary ([SEQ ID NO:8]) amplification primers, respectively, permitted directional cloning of an ~2500 bp PCR product into the commercially available *E.coli* expression plasmid pQE30 (QiaGen, ampicillin resistant) such that a mature BASB027 protein could be expressed as a fusion protein containing a (His)₆ affinity chromatography tag at the N-terminus. The BASB027 PCR product was purified from the amplification reaction using silica gel-based spin columns (QiaGen) according to the manufacturers instructions. To produce the required *Bam*HI and *Sal*I termini necessary for cloning, purified PCR product was sequentially digested to completion with *Bam*HI and *Sal*I restriction enzymes as recommended by the manufacturer (Life Technologies). Following the first restriction digestion, the PCR product was purified via spin column as above to remove salts and eluted in sterile water prior to the second enzyme digestion. The digested DNA fragment was again purified using silica gel-based spin columns prior to ligation with the pQE30 plasmid.

B: Production of Expression Vector.

To prepare the expression plasmid pQE30 for ligation, it was similarly digested to completion with both *Bam*HI and *Sal*I and then treated with calf intestinal phosphatase (CIP, ~0.02 units / pmole of 5' end, Life Technologies) as directed by the manufacturer to prevent self ligation. An approximately 5-fold molar excess of the digested fragment to the prepared vector was used to program the ligation reaction. A standard ~20 µl ligation reaction (~16°C, ~16 hours), using methods well known in the art, was performed using T4 DNA ligase (~2.0 units / reaction, Life Technologies). An aliquot of the ligation (~5 µl) was used to transform electro-competent M15(pREP4) cells according to methods well known in the art. Following a ~2-3 hour outgrowth period at 37°C in ~1.0 ml of LB broth, transformed cells were plated on LB agar plates

containing kanamycin (50 µg/ml) and ampicillin (100 µg/ml). Both antibiotics were included in the selection media to ensure that all transformed cells carried both the pREP4 plasmid (KnR), which carries the *lacIq* gene necessary for the repression of expression for IPTG-inducible expression of proteins on pQE30, and the pQE30-BASB027 plasmid (ApR). Plates were incubated overnight at 37°C for ~16 hours. Individual KnR / ApR colonies were picked with sterile toothpicks and used to "patch" inoculate fresh LB KnR / ApR plates as well as a ~1.0 ml LB KnR / ApR broth culture. Both the patch plates and the broth culture were incubated overnight at 37°C in either a standard incubator (plates) or a shaking water bath.

A whole cell-based PCR analysis was employed to verify that transformants contained the BASB027 DNA insert. Here, the ~1.0 ml overnight LB Kn / Ap broth culture was transferred to a 1.5 ml polypropylene tube and the cells collected by centrifugation in a Beckmann microcentrifuge (~3 min., room temperature, ~12,000 X g). The cell pellet was suspended in ~200µl of sterile water and a ~10µl aliquot used to program a ~50µl final volume PCR reaction containing both BASB027 forward and reverse amplification primers. Final concentrations of the PCR reaction components were essentially the same as those specified in example 2 except ~5.0 units of *Taq* polymerase was used. The initial 95°C denaturation step was increased to 3 minutes to ensure thermal disruption of the bacterial cells and liberation of plasmid DNA. An ABI Model 9700 thermal cycler and a 32 cycle, three-step thermal amplification profile, i.e. 95°C, 45sec; 55-58°C, 45sec, 72°C, 1min., were used to amplify the BASB027 PCR fragment from the lysed transformant samples. Following thermal amplification, a ~20µl aliquot of the reaction was analyzed by agarose gel electrophoresis (0.8 % agarose in a Tris-acetate-EDTA (TAE) buffer). DNA fragments were visualized by UV illumination after gel electrophoresis and ethidium bromide staining. A DNA molecular size standard (1 Kb ladder, Life Technologies) was electrophoresed in parallel with the test samples and was used to estimate the size of the PCR products. Transformants that produced the expected ~ 2500 bp PCR product were identified as strains containing a BASB027

expression construct. Expression plasmid containing strains were then analyzed for the inducible expression of recombinant BASB027.

C: Expression Analysis of PCR-Positive Transformants.

For each PCR-positive transformant identified above, ~5.0 ml of LB broth containing kanamycin (50 µg/ml) and ampicillin (100 µg/ml) was inoculated with cells from the patch plate and grown overnight at 37 °C with shaking (~250 rpm). An aliquot of the overnight seed culture (~1.0 ml) was inoculated into a 125 ml erlenmeyer flask containing ~25 ml of LB Kn / Ap broth and grown at 37 °C with shaking (~250 rpm) until the culture turbidity reached O.D.600 of ~0.5, i.e. mid-log phase (usually about 1.5 - 2.0 hours). At this time approximately half of the culture (~12.5 ml) was transferred to a second 125 ml flask and expression of recombinant BASB027 protein induced by the addition of IPTG (1.0 M stock prepared in sterile water, Sigma) to a final concentration of 1.0 mM. Incubation of both the IPTG-induced and non-induced cultures continued for an additional ~4 hours at 37 °C with shaking. Samples (~1.0 ml) of both induced and non-induced cultures were removed after the induction period and the cells collected by centrifugation in a microcentrifuge at room temperature for ~3 minutes. Individual cell pellets were suspended in ~50µl of sterile water, then mixed with an equal volume of 2X Laemelli SDS-PAGE sample buffer containing 2-mercaptoethanol, and placed in boiling water bath for ~3 min to denature protein. Equal volumes (~15µl) of both the crude IPTG-induced and the non-induced cell lysates were loaded onto duplicate 12% Tris/glycine polyacrylamide gel (1 mm thick Mini-gels, Novex). The induced and non-induced lysate samples were electrophoresed together with prestained molecular weight markers (SeeBlue, Novex) under conventional conditions using a standard SDS/Tris/glycine running buffer (BioRad). Following electrophoresis, one gel was stained with commassie brilliant blue R250 (BioRad) and then destained to visualize novel BASB027 IPTG-inducible protein(s). The second gel was electroblotted onto a PVDF membrane (0.45 micron pore size, Novex) for ~2 hrs at 4 °C using a BioRad Mini-Protean II blotting apparatus and Towbin's methanol (20 %) transfer

buffer. Blocking of the membrane and antibody incubations were performed according to methods well known in the art. A monoclonal anti-RGS (His)₃ antibody, followed by a second rabbit anti-mouse antibody conjugated to HRP (QiaGen), was used to confirm the expression and identity of the BASB027 recombinant protein.

Visualization of the anti-His antibody reactive pattern was achieved using either an ABT insoluble substrate or using Hyperfilm with the Amersham ECL chemiluminescence system.

D: Sequence Confirmation.

To further verify that the IPTG-inducible recombinant BASB027 protein being expressed is in the correct open reading frame and not a spurious molecule arising from a cloning artifact (i.e. a frame-shift), the DNA sequence of the cloned insert was determined. The DNA sequence for the *M. catarrhalis* BASB027 gene was obtained from one strand using conventional asymmetric PCR cycle sequencing methodologies (ABI Prism Dye-Terminator Cycle Sequencing, Perkin-Elmer). Sequencing reactions were programmed with undigested expression plasmid DNA (~0.5µg/rxn) as a template and appropriate pQE30 vector-specific and ORF-specific sequencing primers (~3.5 pmol/rxn). In addition to the template and sequencing primer, each sequencing reaction (~20µl) contained the four different dNTPs (i.e. A,G,C, and T) and the four corresponding ddNTPs (i.e. ddA, ddG, ddC, and ddT) terminator nucleotides; with each terminator being conjugated to one of the four fluorescent dyes, Joe, Tam, Rox, or Fam. Single strand sequencing elongation products were terminated at random positions along the template by the incorporation of the dye-labelled ddNTP terminators. Fluorescent dye-labelled termination products were purified using microcentrifuge size-exclusion chromatography columns (Princeton Genetics), dried under vacuum, suspended in a Template Resuspension Buffer (Perkin-Elmer) for capillary electrophoresis or deionized formamide for PAGE. denatured at 95°C for ~5 min. and analyzed by high resolution capillary electrophoresis (ABI 310 Automated DNA Sequenator, Perkin-Elmer) or high resolution PAGE (ABI 377 Automated DNA

Sequenator) as recommended by the manufacturer. DNA sequence data produced from individual reactions were collected and the relative fluorescent peak intensities analyzed automatically on a PowerMAC computer using ABI Sequence Analysis Software (Perkin-Elmer). Individually autoanalyzed DNA sequences were edited manually for accuracy before being merged into a consensus single strand sequence "string" using AutoAssembler software (Perkin-Elmer). Sequencing determined that the expression plasmid contained the correct sequence in the correct open reading frame.

Example 4 : Production of Recombinant BASB027

Bacterial strain

A recombinant expression strain of *E. coli* M15 (pREP4) containing a plasmid (pQE30) encoding BASB027 from *M. catarrhalis*. was used to produce cell mass for purification of recombinant protein. The expression strain was cultivated on LB agar plates containing 50µg/ml kanamycin ("Kn") and 100µg/ml ampicillin ("Ap") to ensure both the pREP4 lacIq control plasmid and the pQE30-BASB027 expression construct were both maintained. For cryopreservation at -80 °C, the strain was propagated in LB broth containing the same concentration of antibiotics then mixed with an equal volume of LB broth containing 30% (w/v) glycerol.

Media

The fermentation medium used for the production of recombinant protein consisted of 2X YT broth (Difco) containing 50µg/ml Kn and 100µg/ml Ap. Antifoam was added to medium for the fermentor at 0.25 ml/L (Antifoam 204, Sigma). To induce expression of the BASB027 recombinant protein, IPTG (Isopropyl β-D-Thiogalactopyranoside) was added to the fermentor (1 mM, final).

Fermentation

A 500-ml erlenmeyer seed flask, containing 50ml working volume, was inoculated with 0.3 ml of rapidly thawed frozen culture, or several colonies from a selective agar plate culture, and incubated for approximately 12 hours at $37 \pm 1^\circ\text{C}$ on a shaking platform at 150rpm (Innova 2100, New Brunswick Scientific). This seed culture was then used to inoculate a 5-L working volume fermentor containing 2X YT broth and both Kn and Ap antibiotics. The fermentor (Bioflo 3000, New Brunswick Scientific) was operated at $37 \pm 1^\circ\text{C}$, 0.2 - 0.4 VVM air sparge, 250 rpm in Rushton impellers. The pH was not controlled in either the flask seed culture or the fermentor. During fermentation, the pH ranged 6.5 to 7.3 in the fermentor. IPTG (1.0 M stock, prepared in sterile water) was added to the fermentor when the culture reached mid-log of growth (~ 0.7 O.D.600 units). Cells were induced for 2 - 4 hours then harvested by centrifugation using either a 28RS Heraeus (Sepatech) or RC5C superspeed centrifuge (Sorvall Instruments). Cell paste was stored at -20°C until processed.

Purification

Chemicals and Materials

Imidazole, guanidine hydrochloride, Tris (hydroxymethyl), and EDTA (ethylene-diamine tetraacetic acid) biotechnology grade or better were all obtained from Ameresco Chemical, Solon, Ohio. Triton X-100 (t-Octylphenoxypolyethoxy-ethanol), sodium phosphate, monobasic, and Urea were reagent grade or better and obtained from Sigma Chemical Company, St. Louis, Missouri. Glacial acetic acid and hydrochloric acid were obtained from Mallinckrodt Baker Inc., Phillipsburg, New Jersey. Methanol was obtained from Fisher Scientific, Fairlawn, New Jersey. Pefabloc®SC (4-(2-Aminoethyl)-benzenesulfonylfluoride), Complete protease inhibitor cocktail tablets, and PMSF (phenylmethyl-sulfonylfluoride) were obtained from Roche Diagnostics Corporation, Indianapolis, Indiana. Bestatin, Pepstatin A, and E-64 protease inhibitor were obtained from Calbiochem, LaJolla, California. Dulbecco's Phosphate Buffered Saline(1x PBS) was obtained from Quality Biological, Inc., Gaithersburg, Maryland. Dulbecco's Phosphate Buffered Saline (10x PBS) was obtained from BioWhittaker,

Walkersville, Maryland. Penta-His Antibody, BSA free was obtained from QiaGen, Valencia, California. Peroxidase-conjugated AffiniPure Goat Anti-mouse IgG was obtained from Jackson Immuno Research, West Grove, Penn. AEC single solution was obtained from Zymed, South San Francisco, California. All other chemicals were reagent grade or better.

Ni-NTA Superflow resin was obtained from QiaGen Inc., Valencia, California. Precast Tris-Glycine 4-20% and 10-20% polyacrylamide gels, all running buffers and solutions, SeeBlue Pre-Stained Standards, MultiMark Multi-Colored Standards and PVDF transfer membranes were obtained from Novex, San Diego, California. SDS-PAGE Silver Stain kits were obtained from Daiichi Pure Chemicals Company Limited, Tokyo, Japan.

Coomassie Stain Solution was obtained from Bio-Rad Laboratories, Hercules, California. Acrodisc® PF 0.2 µm syringe filters were obtained from Pall Gelman Sciences, Ann Arbor, Michigan. GD/X 25mm disposable syringe filters were obtained from Whatman Inc., Clifton, New Jersey. Dialysis tubing 8,000 MWCO was obtained from BioDesign Inc. Od New York, Carmal New York. BCA Protein Assay Reagents and Snake Skin dialysis tubing 3,500 MWCO were obtained from Pierce Chemical Co., Rockford, Illinois.

Extraction Protocol

Cell paste was thawed at room temperature for 30 to 60 minutes. Five to six grams of material was weighed out into a 50 ml disposable centrifuge tube. To this five mls/gram of Guanidine hydrochloride (Gu-HCl) buffer was added (6 M Guanidine hydrochloride, 100 mM Sodium phosphate, monobasic, 10 mM Tris and 0.05 % Triton X-100, pH 8.0). Cell paste was resuspended using a PRO300D proscientific homogenizer, at 3/4 power for one minute. The extraction mixture was then placed at room temperature with gentle agitation for 60 to 90 minutes. After 60 to 90 minutes the extraction mixture was centrifuged at 15,800 x g for 15 minutes (Sorvall RC5C centrifuge, 11,500 rpm). The supernatant (S1) was decanted and saved for additional purification. The pellet (P1) was saved for analysis.

Binding of BASB027 to Nickel-NTA Resin

To the S1 three to four mls of Ni-NTA resin is added. This is then placed at room temperature with gentle agitation for one hour. After one hour the S1/Ni-NTA is packed into an XK16 Pharmacia column. The column is then washed with 1 M Gu-HCl buffer (1 M Guanidine hydrochloride, 100mM Sodium phosphate, monobasic, 10 mM Tris and 0.05% Triton X-100, pH 8.0). This is then followed by a wash with phosphate buffer (100mM Sodium phosphate, monobasic, 10 mM Tris and 0.05% Triton X-100, pH 6.3). The protein is then eluted from the column with a 250 mM imidazole buffer (250 mM imidazole, 100mM Sodium phosphate, monobasic, 10 mM Tris and 0.05% Triton X-100, pH 5.9).

Final Formulation

BASB027 was formulated by dialysis overnight against, three changes of 0.1 % Triton X-100 and 1x PBS, pH 7.4, to remove residual Gu-HCl and imidazole. The purified protein was characterized and used to produce antibodies as described below.

Biochemical Characterizations

SDS-PAGE and Western Blot Analysis

The recombinant purified protein was resolved on 4-20 % polyacrylamide gels and electrophoretically transferred to PVDF membranes at 100 V for 1 hour as previously described (Thebaine et al. 1979, Proc. Natl. Acad. Sci. USA 76:4350-4354). The PVDF membranes were then pretreated with 25 ml of Dulbecco's phosphate buffered saline containing 5 % non-fat dry milk. All subsequent incubations were carried out using this pretreatment buffer.

PVDF membranes were incubated with 25 ml of a 1:500 dilution of preimmune serum or rabbit anti-His immune serum for 1 hour at room temperature. PVDF membranes were then washed twice with wash buffer (20 mM Tris buffer, pH 7.5, containing 150

mM sodium chloride and 0.05 % Tween-20). PVDF membranes were incubated with 25 ml of a 1:5000 dilution of peroxidase-labeled goat anti-rabbit IgG (Jackson ImmunoResearch Laboratories, West Grove, PA) for 30 minutes at room temperature. PVDF membranes were then washed 4 times with wash buffer, and were developed with 3-amino-9-ethylcarbazole and urea peroxide as supplied by Zymed (San Francisco, CA) for 10 minutes each.

The results of an SDS-PAGE (Figure 4) show a protein about 95 kDa that is reactive to an anti-RGS(His) antibody by western blots (Figure 5) of the SDS-PAGE.

Protein Sequencing

Amino terminal amino acid sequencing of the purified protein was performed to confirm the production of the correct recombinant protein using well defined chemical protocols on Hewlett-Packard model G1000A sequencer with a model 1090 LC and a Hewlett-Packard model 241 sequencer with a model 1100 LC.

Example 5 : Production of Antisera to Recombinant BASB027

Polyvalent antisera directed against the BASB027 protein were generated by vaccinating two rabbits with the purified recombinant BASB027 protein. Each animal is given a total of three immunizations intramuscularly (i.m.) of about 20µg BASB027 protein per injection (beginning with complete Freund's adjuvant and followed with incomplete Freund's adjuvant) at approximately 21 day intervals. Animals were bled prior to the first immunization ("pre-bleed") and on days 35 and 57.

Anti-BASB027 protein titres were measured by an ELISA using purified recombinant BASB027 protein (0.5 µg/well). The titre is defined as the highest dilution equal to or

greater than 0.1 as calculated with the following equation: average OD of two test samples of antisera - the average OD of two test samples of buffer.

The antisera were used as the first antibody to identify the protein in a western blot as described in example 4 above. The western-blot shows the presence of anti-BASB027 antibody in the sera of immunized animals (Figure 6).

Example 6 : Immunological Characterisation

Western Blot Analysis

Several strains of *M. catarrhalis* were grown on chocolate agar plates for 48 hours at 35°C in 5% CO₂. Several colonies were used to inoculate 25ml of Muller Hinton broth in a 250 ml flask. Cultures were grown overnight and collected by centrifugation. Cells were then solubilized by suspending 30µg of cells in 150µl of PAGE sample buffer (360 mM Tris buffer, pH 8.8, containing 4% sodium dodecylsulfate and 20% glycerol), and incubating the suspension at 100°C for 5 minutes. The solubilized cells were resolved on 4-20% polyacrylamide gels and the separated proteins were electrophoretically transferred to PVDF membranes at 100V for 1hour as previously described (Thebaine et al. 1979, Proc. Natl. Acad. Sci. USA 76:4350-4354). The PVDF membranes were then pretreated with 25 ml of Dulbecco's phosphate buffered saline containing 5 % non-fat dry milk. All subsequent incubations were carried out using this pretreatment buffer.

PVDF membranes were incubated with 25ml of a 1:500 dilution of preimmune serum or rabbit immune serum for 1hour at room temperature. PVDF membranes were then washed twice with wash buffer (20 mM Tris buffer, pH 7.5, containing 150 mM sodium chloride and 0.05% Tween-20). PVDF membranes were incubated with 25ml of a 1:5000 dilution of peroxidase-labeled goat anti-rabbit IgG (Jackson ImmunoResearch Laboratories, West Grove, PA) for 30 minutes at room temperature. PVDF membranes were then washed 4 times with wash buffer, and were developed with 3-amino-9-

ethylcarbazole and urea peroxide as supplied by Zymed (San Francisco, CA) for 10 minutes each.

A protein of about 95 kDa (corresponding to BASB027 expected molecular weight) that is reactive with the antisera is detected in all *Moraxella* strains (Figure 7).

Bactericidal Activity

Complement-mediated cytotoxic activity of anti-BASB027 antibodies was examined to determine the vaccine potential of BASB027 polypeptide. Antiserum was prepared as described above. The activities of the pre-immune serum and the anti- BASB027 antiserum in mediating complement killing of *M. catarrhalis* were examined using the "Serum Bactericidal Test" described by Zollinger et al. (Immune Responses to *Neisseria meningitis*, in Manual of Clinical Laboratory Immunology, 3rd ed., pg 347-349), except that cells of *M. catarrhalis* strains or cultivars were used instead of *Neisseria meningitis* cells.

The bactericidal titer of rabbit antiserum (50% killing of homologous strain) was <1:8 (pre-immune) and >1:128 (immune).

Example 7 : Presence of Antibody to BASB027 in Human Convalescent Sera

Western blot analysis of purified recombinant BASB027 were performed as described in Example 4 and 6 above, except that a pool of human sera from children infected by *M. catarrhalis* was used as the first antibody preparation. Results show that antisera from naturally infected individuals react to the purified recombinant.

Example 8 : Production of BASB027 peptides, Antisera and Reactivity Thereof

Two short amino acid BASB027 specific peptides, having the sequences of CYAKPLNKKQNDQTD (SEQ ID NO:9) and YLTARRGQQTTLGEVVC (SEQ ID NO:10) were produced in the laboratory using generally well known methods. These peptides coupled to KLH were used to produce antibodies in 12 weeks old Specific Pathogen Free New-Zealand female rabbits. Rabbits received 4 injections at approximately 3 weeks intervals of 200 µg of peptide-KLH in complete (1st injection) or incomplete (2nd, 3rd and 4th injections) Freund's adjuvant. Animals were bled prior to the first immunization and one month after the 4th injection.

Anti-peptide mid-point titres were measured by an ELISA using free peptides. Anti-peptide Mid-point titres one month after the 4th immunization were superior to 15000. Western blots of purified recombinant BASB027, using anti-peptide antibodies as the first antibody, were prepared as described in Example 4 and 6. The results are presented in Figure 8.

Deposited materials

A deposit containing a *Moraxella catarrhalis* Catlin strain has been deposited with the American Type Culture Collection (herein "ATCC") on June 21, 1997 and assigned deposit number 43617. The deposit was described as *Branhamella catarrhalis* (Frosch and Kolle) and is a freeze-dried, 1.5-2.9 kb insert library constructed from *M. catarrhalis* isolate obtained from a transtracheal aspirate of a coal miner with chronic bronchitis. The deposit is described in Antimicrob. Agents Chemother. 21: 506-508 (1982).

The *Moraxella catarrhalis* strain deposit is referred to herein as "the deposited strain" or as "the DNA of the deposited strain."

The deposited strain contains a full length BASB027 gene.

A deposit of the vector pMC-D15 consisting of *Moraxella catarrhalis* DNA inserted in pQE30 has been deposited with the American Type Culture Collection (ATCC) on February 12 1999 and assigned deposit number 207105.

The sequence of the polynucleotides contained in the deposited strain / clone, as well as the amino acid sequence of any polypeptide encoded thereby, are controlling in the event of any conflict with any description of sequences herein.

The deposit of the deposited strains have been made under the terms of the Budapest Treaty on the International Recognition of the Deposit of Micro-organisms for Purposes of Patent Procedure. The deposited strains will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. The deposited strains are provided merely as convenience to those of skill in the art and are not an admission that a deposit is required for enablement, such as that required under 35 U.S.C. §112.

INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page <u>65</u> , line <u>1-28</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution AMERICAN TYPE CULTURE COLLECTION	
Address of depositary institution (including postal code and country) 10801 UNIVERSITY BLVD, MANASSAS, VIRGINIA 20110-2209, UNITED STATES OF AMERICA	
Date of deposit 21 June 1997 (21.06.97) & 12 February 1999 (12.02.99)	Accession Number 43617 & 207105
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
In respect of those designations where a European Patent is sought, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European Patent or until the date on which the application has been refused or withdrawn, only by issue of such a sample to an expert nominated by the person requesting the sample.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g. "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer R.L.R. Pether	Authorized officer

01-08-2000

BM45324

CLAIMS:

1. An isolated polypeptide comprising an amino acid sequence which has at least 85% identity to the amino acid sequence selected from the group consisting of: SEQ ID NO:2 and SEQ ID NO:4, over the entire length of SEQ ID NO:2 or SEQ ID NO:4 respectively.
2. An isolated polypeptide as claimed in claim 1 in which the amino acid sequence has at least 95% identity to the amino acid sequence selected from the group consisting of: SEQ ID NO:2 and SEQ ID NO:4, over the entire length of SEQ ID NO:2 or SEQ ID NO:4 respectively.
3. The polypeptide as claimed in claim 1 comprising the amino acid sequence selected from the group consisting of: SEQ ID NO:2 and SEQ ID NO:4.
4. An isolated polypeptide having the amino acid sequence selected from the group consisting of SEQ ID NO:2 or SEQ ID NO:4.
5. An immunogenic fragment of the polypeptide as claimed in any one of claims 1 to 4 in which the immunogenic fragment is capable of raising an immune response (if necessary when coupled to a carrier) which recognises the polypeptide of SEQ ID NO:2 or SEQ ID NO:4.
6. A polypeptide as claimed in any of claims 1 to 5 wherein said polypeptide is part of a larger fusion protein.
7. An isolated polynucleotide encoding a polypeptide as claimed in any of claims 1 to 6.
8. An isolated polynucleotide comprising a nucleotide sequence encoding a polypeptide that has at least 85% identity to the amino acid sequence of SEQ ID NO:2 or 4 over the entire length of SEQ ID NO:2 or 4 respectively; or a nucleotide sequence complementary to said isolated polynucleotide.

01-08-2000

BM45324

9. An isolated polynucleotide comprising a nucleotide sequence that has at least 85% identity to a nucleotide sequence encoding a polypeptide of SEQ ID NO:2 or 4 over the entire coding region; or a nucleotide sequence complementary to said isolated polynucleotide.

5

10. An isolated polynucleotide which comprises a nucleotide sequence which has at least 85% identity to that of SEQ ID NO:1 or 3 over the entire length of SEQ ID NO:1 or 3 respectively; or a nucleotide sequence complementary to said isolated polynucleotide.

10 11. The isolated polynucleotide as claimed in any one of claims 7 to 10 in which the identity is at least 95% to SEQ ID NO:1 or 3.

12. An isolated polynucleotide comprising a nucleotide sequence encoding the polypeptide of SEQ ID NO:2 or SEQ ID NO:4.

15

13. An isolated polynucleotide comprising the polynucleotide of SEQ ID NO:1 or SEQ ID NO:3.

20

14. An isolated polynucleotide comprising a nucleotide sequence encoding the polypeptide of SEQ ID NO:2, SEQ ID NO:4 obtainable by screening an appropriate library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO:1 or SEQ ID NO:3 or a fragment thereof.

25

15. An expression vector or a live microorganism comprising an isolated recombinant polynucleotide according to any one of claims 7 - 14.

30

16. A host cell comprising the expression vector of claim 15 expressing an isolated polypeptide comprising an amino acid sequence that has at least 85% identity to the amino acid sequence selected from the group consisting of: SEQ ID NO:2 and SEQ ID NO:4, or a membrane of the host cell comprising the expressed polypeptide.

01-08-2000

BM45324

17. A process for producing a polypeptide of claims 1 to 6 comprising culturing a host cell of claim 16 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture medium.

5 18. A process for expressing a polynucleotide of any one of claims 7 - 14 comprising transforming a host cell with the expression vector comprising at least one of said polynucleotides and culturing said host cell under conditions sufficient for expression of any one of said polynucleotides.

10 19. A vaccine composition comprising an effective amount of the polypeptide of any one of claims 1 to 6 and a pharmaceutically acceptable carrier.

20. A vaccine composition comprising an effective amount of the polynucleotide of any one of claims 7 to 14 and a pharmaceutically effective carrier.

15 21. The vaccine composition according to either one of claims 19 or 20 wherein said composition comprises at least one other *Neisseria meningitidis* antigen.

20 22. An antibody immunospecific for the polypeptide or immunological fragment as claimed in any one of claims 1 to 6.

25 23. A method of diagnosing a *Neisseria meningitidis* infection, comprising identifying a polypeptide as claimed in any one of claims 1 - 6, or an antibody that is immunospecific for said polypeptide, present within a biological sample from an animal suspected of having such an infection.

24. Use of a composition comprising an immunologically effective amount of a polypeptide as claimed in any one of claims 1 - 6 in the preparation of a medicament for use in generating an immune response in an animal.

30

01-08-2000

BM45324

25. Use of a composition comprising an immunologically effective amount of a polynucleotide as claimed in any one of claims 7 - 14 in the preparation of a medicament for use in generating an immune response in an animal.


- 5 26. A therapeutic composition useful in treating humans with *Neisseria meningitidis* comprising at least one antibody directed against the polypeptide of claims 1 - 6 and a suitable pharmaceutical carrier.

10


10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500 1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770 1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800 1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1819 1820 1821 1822 1823 1824 1825 1826 1827 1828 1829 1830 1831 1832 1833 1834 1835 1836 1837 1838 1839 1840 1841 1842 1843 1844 1845 1846 1847 1848 1849 1850 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860 1861 1862 1863 1864 1865 1866 1867 1868 1869 1870 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890 1891 1892 1893 1894 1895 1896 1897 1898 1899 1900 1901 1902 1903 1904 1905 1906 1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590 2591 25

1/26


Figure 1 : Photographs of the electrophoresis gels obtained in the RFLP analysis of BASB027




2/26




3/26




5/26



6/26




7126



SUBSTITUTE SHEET (RULE 26)

[illegible]

8/26




09/701711


WO 99/63093

PCT/EP99/03822


9/26




10/26



11/26





13/26

Figure 2 : Alignment of the BASB027 polynucleotide sequences.**Identity to SeqID No:1 is indicated by a dot.**

```

                *      20      *      40      *
Seqid1 : ATGCGTAATTCATATTTTAAAGGTTTTCAGGTCAGTGCAATGACAATGGC : 50
Seqid3 : ..... : 50

                60      *      80      *      100
Seqid1 : TGTGATGATGGTAATGTCAACTCATGCACAAGCGGCGGATTTTATGGCAA : 100
Seqid3 : ..... : 100

                *      120      *      140      *
Seqid1 : ATGACATTACCATCACAGGACTACAGCGAGTGACCATTGAAAGCTTACAA : 150
Seqid3 : .....G..... : 150

                160      *      180      *      200
Seqid1 : AGCGTGCTGCCGTTTCGCTTGGGTCAAGTGGTGAGCGAAAACCAAGTTGGC : 200
Seqid3 : .....GCA..... : 200

                *      220      *      240      *
Seqid1 : TGATGGTGTCAAAGCACTTTATGCAACAGGCAATTTTTCAGATGTGCAAG : 250
Seqid3 : ..... : 250

                260      *      280      *      300
Seqid1 : TCTATCATCAAGAAGGGCGTATCATCTATCAGGTAACCGAAAGGCCGTTA : 300
Seqid3 : ..... : 300

                *      320      *      340      *
Seqid1 : ATCGCTGAGATTAATTTTGAGGGCAATCGCTTAATTCCAAAGAAGGTCT : 350
Seqid3 : ..... : 350

```

14 / 26

360 * 380 * 400
Seqid1 : ACAAGAAGGGCTAAAAAATGCTGGCTTAGCTGTGGGTCAACCACTAAAAC : 400
Seqid3 : : 400

* 420 * 440 *
Seqid1 : AAGCCACAGTACAGATGATCGAAACCGAGCTTACCAATCAATATATATCA : 450
Seqid3 : : 450

460 * 480 * 500
Seqid1 : CAAGGCTATTATAATACCGAAATTACTGTCAAACAGACGATGCTTGATGG : 500
Seqid3 : : 500

* 520 * 540 *
Seqid1 : TAATCGTGTTAAGCTTGATATGACCTTTGCTGAAGGTAAACCTGCACGGG : 550
Seqid3 : : 550

560 * 580 * 600
Seqid1 : TGGTTGATATTAATATCATTGGCAATCAGCATTTTAGCGATGCAGATTTG : 600
Seqid3 : : 600

* 620 * 640 *
Seqid1 : ATTGATGTGCTTGCGATTAAGGATAATAAAATCAATCCACTGTCTAAAGC : 650
Seqid3 : : 650

660 * 680 * 700
Seqid1 : TGACCGTTATACTCAAGAAAAGCTGGTGACCAGTTTAGAGAATTTGCGTG : 700
Seqid3 : : 700

* 720 * 740 *
Seqid1 : CTAAATATCTCAATGCAGGGTTTGTGCGTTTTGAGATTAAAGATGCTAAG : 750

15/26

Seqid3 : : 750

760 * 780 * 800
Seqid1 : CTTAATATTAATGAAGATAAAAACCGTATCTTTGTTGAGATTCATTGCA : 800
Seqid3 : : 800

* 820 * 840 *
Seqid1 : TGAAGGTGAGCAATATCGCTTTGGACAGACACAGTTTTTGGGTAATTTAA : 850
Seqid3 : : 850

860 * 880 * 900
Seqid1 : CTTATACTCAAGCAGAACTTGAGGCACTGCTTAAATTCAAAGCAGAAGAA : 900
Seqid3 : : 900

* 920 * 940 *
Seqid1 : GGGTTTTCAACAAGCCATGCTTGAGCAAACAACAACAATATCAGTACCAA : 950
Seqid3 : : 950

960 * 980 * 1000
Seqid1 : ATTTGGTGACGATGGCTATTATTATGCTCAAATCCGTCCTGTAAACACGCA : 1000
Seqid3 : : 1000

* 1020 * 1040 *
Seqid1 : TTAATGATGAAAGTCGTACGGTTGATGTGGAATATTATATTGACCCTGTA : 1050
Seqid3 : : 1050

1060 * 1080 * 1100
Seqid1 : CACCCTGTCTATGTACGCCGTATTAATTTTACAGGTAACTTAAGACCCA : 1100
Seqid3 : : 1100

16/26

 * 1120 * 1140 *
Seqid1 : AGATGAAGTACTCCGTCGTGAGATGCGACAACTTGAAGGTGCGTTGGCAT : 1150
Seqid3 : : 1150

 1160 * 1180 * 1200
Seqid1 : CTAATCAAAAAATCCAGCTGTCTCGTGACGCTTGATGCGGACTGGGTTT : 1200
Seqid3 : : 1200

 * 1220 * 1240 *
Seqid1 : TTAAACATGTTACCGTTGATACTCGTCCAGTACCCAACTCACCTGATCA : 1250
Seqid3 : : 1250

 1260 * 1280 * 1300
Seqid1 : GGTGATGTAAATTTTGTGGTTGAAGAACAACCTTCAGGATCATCAACCA : 1300
Seqid3 : : 1300

 * 1320 * 1340 *
Seqid1 : TCGCAGCAGGCTACTCTCAAAGTGGTGGTGTAACCTTTTCAATTTGATGTT : 1350
Seqid3 : : 1350

 1360 * 1380 * 1400
Seqid1 : TCTCAAATAAATTTATGGGTACAGGTAAGCACGTCAATGCTTCGTTTTTC : 1400
Seqid3 : : 1400

 * 1420 * 1440 *
Seqid1 : TCGCTCTGAGACCCGTGAGGTGTATAGTTTGGGTATGACCAACCCATACT : 1450
Seqid3 : : 1450

 1460 * 1480 * 1500
Seqid1 : TTACCGTAAATGGCGTCTCGCAAAGCTTGAGTGGCTACTATCGTAAAACC : 1500
Seqid3 : : 1500

17/26

 * 1520 * 1540 *
Seqid1 : AAGTATGATAACAAGAACATTAGTAATTATGTACTTGATTCTTATGGTGG : 1550
Seqid3 : : 1550

 1560 * 1580 * 1600
Seqid1 : CTCATTAAGCTATGGATATCCAATTGATGAAAATCAACGCATAAGCTTTG : 1600
Seqid3 : : 1600

 * 1620 * 1640 *
Seqid1 : GTCTGAATGCTGACAATACCAAGCTTCATGGCGGTCGTTTTATGGGCATT : 1650
Seqid3 : : 1650

 1660 * 1680 * 1700
Seqid1 : AGTAATGTCAAGCAGCTGATGGCAGATGGTGGCAAATTCAGTGGATAA : 1700
Seqid3 : : 1700

 * 1720 * 1740 *
Seqid1 : TAATGGCATTCTGATTTTAAGCATGATTACACAACCTACAATGCCATTT : 1750
Seqid3 : : 1750

 1760 * 1780 * 1800
Seqid1 : TGGGGTGGAAATTATTCAAGTCTAGATCGCCCTGTATTTCCAACCCAAGGC : 1800
Seqid3 : : 1800

 * 1820 * 1840 *
Seqid1 : ATGAGTCATTCTGTAGATTTGACGGTTGGTTTTGGTGATAAACTCATCA : 1850
Seqid3 : : 1850

 1860 * 1880 * 1900
Seqid1 : AAAAGTGGTTTATCAAGGCAATATCTATCGCCCATTTATCAAAAAATCAG : 1900

18/26

Seqid3 : : 1900

* 1920 * 1940 *

Seqid1 : TCTTGCGTGGATACGCCAAGTTAGGCTATGGCAATAATTTACCATTTTAT : 1950

Seqid3 : : 1950

1960 * 1980 * 2000

Seqid1 : GAAAATTTCTATGCAGGCGGCTATGGTTCGGTTCGTGGCTATGATCAATC : 2000

Seqid3 : : 2000

* 2020 * 2040 *

Seqid1 : CTCTTTGGGTCCACGCTCACAAGCCTATTTGACAGCTCGTCGTGGTCAAC : 2050

Seqid3 : : 2050

2060 * 2080 * 2100

Seqid1 : AAACCACACTAGGAGAGGTTGTTGGTGGTAATGCTTTGGCAACTTTCGGC : 2100

Seqid3 : : 2100

* 2120 * 2140 *

Seqid1 : AGTGAGCTGATTTTACCTTTGCCATTTAAAGGTGATTGGATAGATCAGGT : 2150

Seqid3 : : 2150

2160 * 2180 * 2200

Seqid1 : GCGTCCAGTGATATTCATTGAGGGCGGTCAGGTTTTTGATACAACAGGTA : 2200

Seqid3 : : 2200

* 2220 * 2240 *

Seqid1 : TGGATAAACAAACCATTGATTTAACCCAATTTAAAGACCCACAAGCAACA : 2250

Seqid3 : : 2250

19/26

2260 * 2280 * 2300
Seqid1 : GCTGAACAAAATGCAAAAGCAGCCATCGCCCGCTACTAACCCCAAGATAA : 2300
Seqid3 : : 2300

* 2320 * 2340 *
Seqid1 : ACAGTTGCGTTATAGTGCTGGTGTTGGTGCAACTTGGTATACGCCCCATTG : 2350
Seqid3 : : 2350

2360 * 2380 * 2400
Seqid1 : GTCCTTTATCTATTAGCTATGCCAAGCCATTGAATAAAAAACAAAATGAT : 2400
Seqid3 : : 2400

* 2420 * 2440
Seqid1 : CAGACCGATACGGTACAGTTCCAGATTGGTAGTGTCTTTTAA : 2442
Seqid3 : : 2442

20/26

Figure 3 : Alignment of the BASB027 polypeptide sequences.

Identity to SeqID No:2 is indicated by a dot.

```

          *          20          *          40          *
Seqid2 : MRNSYFKGFQVSAMTMAVMMVMSTHAQAADFMANDITITGLQRVTTIESLQ : 50
Seqid4 : .....A..... : 50

          60          *          80          *          100
Seqid2 : SVLPFRLGQVVSENQLADGVKALYATGNFSDVQVYHQEGRIIYQVTERPL : 100
Seqid4 : .....A..... : 100

          *          120          *          140          *
Seqid2 : IAEINFEGNRLIPKEGLQEGLKNAGLAVGQPLKQATVQMIETELTNQYIS : 150
Seqid4 : ..... : 150

          160          *          180          *          200
Seqid2 : QGYNTEITVKQTMLDGNRVKLDMTFAEGKPARVVDINIIGNQHFSDADL : 200
Seqid4 : ..... : 200

          *          220          *          240          *
Seqid2 : IDVLAIKDNKINPLSKADRYTQEKLVTSLENLRAKYLNAGFVRFEIKDAK : 250
Seqid4 : ..... : 250

          260          *          280          *          300
Seqid2 : LNINEDKNRIFVEISLHEGEQYRFGQTQFLGNLTYTQAELEALLKFKAEE : 300
Seqid4 : ..... : 300

          *          320          *          340          *
Seqid2 : GFSQAMLEQTTNNISTKFGDDGYYYAQIRPVTRINDESRTVDVEYYIDPV : 350
Seqid4 : ..... : 350

```


21/26

360 * 380 * 400
Seqid2 : HPVYVRRINFTGNFKTQDEVLRREMRQLEGALASNQKIQLSRARLMRTGF : 400
Seqid4 : : 400

* 420 * 440 *
Seqid2 : FKHVTVDTRPVPNSPDQVDVNFVVEEQPSGSSTIAAGYSQSGGVTFQFDV : 450
Seqid4 : : 450

460 * 480 * 500
Seqid2 : SQNNFMGTGKHVNASFSRSETREVYSLGMTNPYFTVNGVSQSLSGYYRKT : 500
Seqid4 : : 500

* 520 * 540 *
Seqid2 : KYDNKNISNYVLDSYGGSLSYGYPIDENQRISFGLNADNTKLHGGRFMGI : 550
Seqid4 : : 550

560 * 580 * 600
Seqid2 : SNVKQLMADGGKIQVDNNGIPDFKHDTTYNAILGWNYSLLDRPVFPTQG : 600
Seqid4 : : 600

* 620 * 640 *
Seqid2 : MSHSVDLTVGFGDKTHQKVYQGNIRPFIKKSVLRGYAKLGYGNNLPFY : 650
Seqid4 : : 650

660 * 680 * 700
Seqid2 : ENFYAGGYGSVRGYDQSSLGPRSQAYLTARRGQQTTLGEVVGGNALATFG : 700
Seqid4 : : 700

* 720 * 740 *
Seqid2 : SELILPLPFKGDWIDQVRPVIFIEGGQVFDTTGMDKQTIDLTQFKDPQAT : 750

22/26

Seqid4 : : 750

760

*

780

*

800

Seqid2 : AEQNAKAANRPLLTDKQLRYSAGVGATWYTPIGPLSISYAKPLNKKQNE : 800

Seqid4 : : 800

*

Seqid2 : QTDTVQFQIGSVF : 813

Seqid4 : : 813

23/26

Figure 4: Coomassie stained SDS-PAGE of purified BASB027 protein.






Figure 5: Western blot with anti-His antibody of purified BASB027 protein.




24/26

Figure 6: Western blot of purified BASB027 protein with the corresponding anti-recombinant protein sera. Panel A: pre-immune serum. Panel B: immune serum.



25/26


Figure 7: Western blot of whole cell lysates of 16 strains of *M. catarrhalis* using pooled sera against the BASB027 protein (sera was diluted 1:2000).



26/26

Figure 8: Western blot of purified recombinant BASB027 with corresponding anti-peptide sera. Lanes 2 and 3 non immune sera. Lanes 1 and 4 immune sera.

MWM 1 2 3 4



WO 99/63093

PCT/EP99/03822

SEQUENCE LISTING

<110> SmithKline Beecham Biologicals

<120> Novel Compounds

<130> BM45324

<160> 10

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 2442

<212> DNA

<213> Bacteria

<400> 1

atgcgtaatt	catatttttaa	agggttttcag	gtcagtgcaa	tgacaatggc	tgatcatgatg	60
gtaatgtcaa	ctcatgcaca	agcggcgcat	tttatggcaa	atgacattac	catcacagga	120
ctacagcgag	tgaccattga	aagcttacaa	agcgtgctgc	cgtttcgctt	gggtcaagtg	180
gtgagcgaaa	accagttggc	tgatgggtgc	aaagcacttt	atgcaacagg	caatttttca	240
gatgtgcaag	tctatcatca	agaagggcgt	atcatctatc	aggtaaccca	aaggccgtta	300
atcgctgaga	ttaattttga	gggcaatcgc	ttaattccaa	aagaaggtct	acaagaaggg	360
ctaaaaaatg	ctggcttagc	tgtgggtcaa	ccactaaaac	aagccacagt	acagatgatc	420
gaaaccgagc	ttaccaatca	atatatatca	caaggctatt	ataataccga	aattactgtc	480
aaacagacga	tgcttgatgg	taatcgtggt	aagcttgata	tgacctttgc	tgaaggtaaa	540
cctgcacggg	tggttgatat	taatatcatt	ggcaatcagc	attttagcga	tgagagattg	600
attgatgtgc	ttgcgattaa	ggataataaa	atcaatccac	tgtctaaagc	tgaccgttat	660
actcaagaaa	agctggtgac	cagtttagag	aatttgcgtg	ctaaatatct	caatgcaggg	720
tttggtgctt	ttgagattaa	agatgctaag	cttaatatta	atgaagataa	aaaccgtatc	780
tttggttgaga	tttcattgca	tgaaggtgag	caatatcgct	ttggacagac	acagtttttg	840
ggtaattttaa	cttatactca	agcagaactt	gaggcactgc	ttaaattcaa	agcagaagaa	900
gggtttttcac	aagccatgct	tgagcaaaaca	acaaacaata	tcagtaccaa	atttggtgac	960
gatggctatt	attatgctca	aatccgtcct	gtaacacgca	ttaatgatga	aagtcgtacg	1020
gttgatgtgg	aatattatat	tgaccctgta	cacctgtctt	atgtacgcgg	tattaatttt	1080
acaggtaact	ttaagaccca	agatgaagta	ctccgtcgtg	agatgcgaca	acttgaagggt	1140
gcgtttggcat	ctaatacaaaa	aatccagctg	tctcgtgcac	gcttgatgcy	gactgggttt	1200
tttaaacatg	ttaccgttga	tactcgtcca	gtaccaactt	cacctgatca	ggttgatgta	1260
aattttgtgg	ttgaagaaca	accttcagga	tcatcaacca	tcgcagcagg	ctactctcaa	1320
agtgggtggg	taacttttca	atttgatggt	tctcaaaaata	actttatggg	tacaggtaag	1380
cacgtcaatg	cttcgttttc	tgcgtctgag	acccgtgagg	tgtatagttt	gggtatgacc	1440
aaccatactt	ttaccgtaaa	tggcgtctcg	caaagcttga	gtggctacta	tcgtaaaacc	1500
aagtatgata	acaagaacat	tagtaattat	gtacttgatt	cttatgggtg	ctcattaagc	1560
tatggatatt	caattgatga	aaatcaacgc	ataagctttg	gtctgaatgc	tgacaatacc	1620
aagcttcatg	gcggtcgttt	tatgggcatt	agtaatgtca	agcagctgat	ggcagatggt	1680
ggcaaaaattc	aagtggataa	taatggcatt	cctgatttta	agcatgatta	cacaacctac	1740
aatgccattt	tgggggtggaa	ttattcaagt	ctagatcgcc	ctgtattttc	aaccgaaggc	1800
atgagtcatt	ctgtagattt	gacgggttgg	tttggtgata	aaactcatca	aaaagtgggt	1860
tatcaaggca	atatctatcg	cccattttatc	aaaaaatcag	tcttgcgtgg	atacgccaag	1920
ttaggctatg	gcaataattt	accattttat	gaaaatttct	atgcaggcgg	ctatggttcg	1980
gttcgtggct	atgatcaatc	ctctttgggt	ccacgctcac	aagcctattt	gacagctcgt	2040
cgtgggtcaac	aaaccacatt	aggagaggtt	gttgggtgta	atgctttggc	aactttcggc	2100
agtgaactga	ttttaccttt	gccattttaa	ggtgattgga	tagatcaggt	gcgtccagtg	2160
atattcattg	agggcggtca	ggtttttgat	acaacaggta	tggataaaca	aaccattgat	2220
tttaacccaat	ttaaagaccc	acaagcaaca	gctgaacaaa	atgcaaaagc	agccaatcgc	2280
ccgctactaa	cccaagataa	acagttgcgt	tatagtgctg	gtgttggtgc	aacttggtat	2340

acgcccattg gtcctttatc tattagctat gccaaagccat tgaataaaaa acaaaatgat 2400
cagaccgata cggtagcagtt ccagattgggt agtgtctttt aa 2442

<210> 2
<211> 813
<212> PRT
<213> Bacteria

<400> 2
Met Arg Asn Ser Tyr Phe Lys Gly Phe Gln Val Ser Ala Met Thr Met
1 5 10 15
Ala Val Met Met Val Met Ser Thr His Ala Gln Ala Ala Asp Phe Met
20 25 30
Ala Asn Asp Ile Thr Ile Thr Gly Leu Gln Arg Val Thr Ile Glu Ser
35 40 45
Leu Gln Ser Val Leu Pro Phe Arg Leu Gly Gln Val Val Ser Glu Asn
50 55 60
Gln Leu Ala Asp Gly Val Lys Ala Leu Tyr Ala Thr Gly Asn Phe Ser
65 70 75 80
Asp Val Gln Val Tyr His Gln Glu Gly Arg Ile Ile Tyr Gln Val Thr
85 90 95
Glu Arg Pro Leu Ile Ala Glu Ile Asn Phe Glu Gly Asn Arg Leu Ile
100 105 110
Pro Lys Glu Gly Leu Gln Glu Gly Leu Lys Asn Ala Gly Leu Ala Val
115 120 125
Gly Gln Pro Leu Lys Gln Ala Thr Val Gln Met Ile Glu Thr Glu Leu
130 135 140
Thr Asn Gln Tyr Ile Ser Gln Gly Tyr Tyr Asn Thr Glu Ile Thr Val
145 150 155 160
Lys Gln Thr Met Leu Asp Gly Asn Arg Val Lys Leu Asp Met Thr Phe
165 170 175
Ala Glu Gly Lys Pro Ala Arg Val Val Asp Ile Asn Ile Ile Gly Asn
180 185 190
Gln His Phe Ser Asp Ala Asp Leu Ile Asp Val Leu Ala Ile Lys Asp
195 200 205
Asn Lys Ile Asn Pro Leu Ser Lys Ala Asp Arg Tyr Thr Gln Glu Lys
210 215 220
Leu Val Thr Ser Leu Glu Asn Leu Arg Ala Lys Tyr Leu Asn Ala Gly
225 230 235 240
Phe Val Arg Phe Glu Ile Lys Asp Ala Lys Leu Asn Ile Asn Glu Asp
245 250 255
Lys Asn Arg Ile Phe Val Glu Ile Ser Leu His Glu Gly Glu Gln Tyr
260 265 270
Arg Phe Gly Gln Thr Gln Phe Leu Gly Asn Leu Thr Tyr Thr Gln Ala
275 280 285
Glu Leu Glu Ala Leu Leu Lys Phe Lys Ala Glu Glu Gly Phe Ser Gln
290 295 300
Ala Met Leu Glu Gln Thr Thr Asn Asn Ile Ser Thr Lys Phe Gly Asp
305 310 315 320
Asp Gly Tyr Tyr Tyr Ala Gln Ile Arg Pro Val Thr Arg Ile Asn Asp
325 330 335
Glu Ser Arg Thr Val Asp Val Glu Tyr Tyr Ile Asp Pro Val His Pro
340 345 350
Val Tyr Val Arg Arg Ile Asn Phe Thr Gly Asn Phe Lys Thr Gln Asp
355 360 365
Glu Val Leu Arg Arg Glu Met Arg Gln Leu Glu Gly Ala Leu Ala Ser
370 375 380
Asn Gln Lys Ile Gln Leu Ser Arg Ala Arg Leu Met Arg Thr Gly Phe


```

385          390          395          400
Phe Lys His Val Thr Val Asp Thr Arg Pro Val Pro Asn Ser Pro Asp
          405          410          415
Gln Val Asp Val Asn Phe Val Val Glu Glu Gln Pro Ser Gly Ser Ser
          420          425          430
Thr Ile Ala Ala Gly Tyr Ser Gln Ser Gly Gly Val Thr Phe Gln Phe
          435          440          445
Asp Val Ser Gln Asn Asn Phe Met Gly Thr Gly Lys His Val Asn Ala
          450          455          460
Ser Phe Ser Arg Ser Glu Trp Arg Glu Val Tyr Ser Leu Gly Met Thr
          465          470          475          480
Asn Pro Tyr Phe Thr Val Asn Gly Val Ser Gln Ser Leu Ser Gly Tyr
          485          490          495
Tyr Arg Lys Thr Lys Tyr Asp Asn Lys Asn Ile Ser Asn Tyr Val Leu
          500          505          510
Asp Ser Tyr Gly Gly Ser Leu Ser Tyr Gly Tyr Pro Ile Asp Glu Asn
          515          520          525
Gln Arg Ile Ser Phe Gly Leu Asn Ala Asp Asn Thr Lys Leu His Gly
          530          535          540
Gly Arg Phe Met Gly Ile Ser Asn Val Lys Gln Leu Met Ala Asp Gly
          545          550          555          560
Gly Lys Ile Gln Val Asp Asn Asn Gly Ile Pro Asp Phe Lys His Asp
          565          570          575
Tyr Thr Thr Tyr Asn Ala Ile Leu Gly Trp Asn Tyr Ser Ser Leu Asp
          580          585          590
Arg Pro Val Phe Pro Thr Gln Gly Met Ser His Ser Val Asp Leu Thr
          595          600          605
Val Gly Phe Gly Asp Lys Thr His Gln Lys Val Val Tyr Gln Gly Asn
          610          615          620
Ile Tyr Arg Pro Phe Ile Lys Lys Ser Val Leu Arg Gly Tyr Ala Lys
          625          630          635          640
Leu Gly Tyr Gly Asn Asn Leu Pro Phe Tyr Glu Asn Phe Tyr Ala Gly
          645          650          655
Gly Tyr Gly Ser Val Arg Gly Tyr Asp Gln Ser Ser Leu Gly Pro Arg
          660          665          670
Ser Gln Ala Tyr Leu Thr Ala Arg Arg Gly Gln Gln Thr Thr Leu Gly
          675          680          685
Glu Val Val Gly Gly Asn Ala Leu Ala Thr Phe Gly Ser Glu Leu Ile
          690          695          700
Leu Pro Leu Pro Phe Lys Gly Asp Trp Ile Asp Gln Val Arg Pro Val
          705          710          715          720
Ile Phe Ile Glu Gly Gly Gln Val Phe Asp Thr Thr Gly Met Asp Lys
          725          730          735
Gln Thr Ile Asp Leu Thr Gln Phe Lys Asp Pro Gln Ala Thr Ala Glu
          740          745          750
Gln Asn Ala Lys Ala Ala Asn Arg Pro Leu Leu Thr Gln Asp Lys Gln
          755          760          765
Leu Arg Tyr Ser Ala Gly Val Gly Ala Thr Trp Tyr Thr Pro Ile Gly
          770          775          780
Pro Leu Ser Ile Ser Tyr Ala Lys Pro Leu Asn Lys Lys Gln Asn Asp
          785          790          795          800
Gln Thr Asp Thr Val Gln Phe Gln Ile Gly Ser Val Phe
          805          810

```

<210> 3
 <211> 2442
 <212> DNA
 <213> Bacteria

<400> 3

```

atgcgtaatt catattttaa aggttttcag gtcagtgcac tgacaatggc tgtcatgatg      60
gtaatgtcaa ctcatgcaca agcggcggat tttatggcaa atgacattgc catcacagga      120
ctacagcgag tgaccattga aagcttacaa agcgtgctgc cgtttcgttt gggccaagtg      180
gtgagcgaa cagagttggc tgatggtgtc aaagcacttt atgcaacagg caatttttca      240
gatgtgcaag tctatcatca agaagggcgt atcatctatc aggtaaccga aaggccgtta      300
atcgctgaga ttaattttga gggcaatcgc ttaattccaa aagaaggctc acaagaaggg      360
ctaaaaaatg ctggccttagc tgtgggtcaa ccactaaaac aagccacagt acagatgatc      420
gaaaccgagc ttaccaatca atatatatca caaggctatt ataataccga aattactgtc      480
aaacagacga tgcttgatgg taatcgtgtt aagcttgata tgacctttgc tgaaggtaaa      540
cctgcacggg tgggttgatat taatatcatt ggcaatcagc attttagcga tgcagatttg      600
attgatgtgc ttgcgattaa ggataataaa atcaatecac tgtctaaagc tgaccgttat      660
actcaagaaa agctgggtgac cagtttagag aatttgcgtg ctaaatatct caatgcaggg      720
tttgtgcgtt ttgagattaa agatgctaag cttaattata atgaagataa aaaccgtatc      780
tttgttgaga ttctattgca tgaagggtgag caatatcgct ttggacagac acagtttttg      840
ggtaatttaa cttatactca agcagaactt gaggcactgc ttaaattcaa agcagaagaa      900
gggttttcac aagccatgct tgagcaaaaca acaacaata tcagtaccaa atttgggtgac      960
gatggctatt attatgctca aatccgtcct gtaacacgca ttaatgatga aagtcgtacg      1020
gttgatgtgg aatattatat tgacctgta caccctgtct atgtacgccg tattaatttt      1080
acaggtaact ttaagaccca agatgaagta ctccgtcgtg agatgcgaca acttgaaggt      1140
gcgttggcat ctaatcaaaa aatccagctg tctcgtgcac gcttgatgcg gactgggttt      1200
tttaaacatg ttaccgttga tactcgtcca gtacccaact cacctgatca ggttgatgta      1260
aattttgtgg ttgaagaaca accttcagga tcatcaacca tcgcagcagg ctactctcaa      1320
agtgggtggg taacttttca atttgatgtt tctcaaaaata accttatggg tacaggtaag      1380
cacgtcaatg cttcgttttc tcgctctgag acccgtgagg tgtatagttt gggtagtacc      1440
aaccatact ttaccgtaaa tggcgtctcg caaagcttga gtggctacta tcgtaaaacc      1500
aagtatgata acaagaacat tagtaattat gtacttgatt cttatgggtg ctcattaagc      1560
tatggatata caattgatga aaatcaacgc ataagctttg gtctgaatgc tgacaatacc      1620
aagcttcatg gcggtcgttt tatgggcatt agtaatgtca agcagctgat ggcagatggg      1680
ggcaaaaattc aagtggataa taatggcatt cctgatttta agcatgatta cacaacctac      1740
aatgccattt tgggggtggaa ttattcaagt ctagatcgcc ctgtatttcc aaccaaggc      1800
atgagtcatt ctgtagattt gacgggtggg tttggtgata aaactcatca aaaagtgggt      1860
tatcaaggca atattctatcg cccatttatc aaaaaatcag tcttgcgtgg atacgccaag      1920
ttaggctatg gcaataattt accattttat gaaaaattct atgcaggcgg ctatggttcg      1980
gttcgtggct atgatcaatc ctctttgggt ccacgctcac aagcctattt gacagctcgt      2040
cgtgggtcaac aaaccacact aggagaggtt gttgggtgga atgctttggc aactttcggc      2100
agtgaagctga ttttaccttt gccattttaa ggtgattgga tagatcaggt gcgtccagtg      2160
atattcattg agggcgggtca ggtttttgat acaacaggta tggataaaca aaccattgat      2220
ttaaccat ttaaagaccc acaagcaaca gctgaacaaa atgcaaaagc agccaatcgc      2280
ccgctactaa cccaagataa acagttgcgt tatagtgtcg gtgttgggtg aacttgggtat      2340
acgcccattg gtcttttatc tattagctat gccaaagccat tgaataaaaa acaaatgat      2400
cagaccgata cggtagagtt ccagattggt agtgtctttt aa      2442

```

<210> 4

<211> 813

<212> PRT

<213> Bacteria

<400> 4

```

Met Arg Asn Ser Tyr Phe Lys Gly Phe Gln Val Ser Ala Met Thr Met
1           5           10           15
Ala Val Met Met Val Met Ser Thr His Ala Gln Ala Ala Asp Phe Met
20           25           30
Ala Asn Asp Ile Ala Ile Thr Gly Leu Gln Arg Val Thr Ile Glu Ser
35           40           45
Leu Gln Ser Val Leu Pro Phe Arg Leu Gly Gln Val Val Ser Glu Ala
50           55           60

```

Gln Leu Ala Asp Gly Val Lys Ala Leu Tyr Ala Thr Gly Asn Phe Ser
 65 70 75 80
 Asp Val Gln Val Tyr His Gln Glu Gly Arg Ile Ile Tyr Gln Val Thr
 85 90 95
 Glu Arg Pro Leu Ile Ala Glu Ile Asn Phe Glu Gly Asn Arg Leu Ile
 100 105 110
 Pro Lys Glu Gly Leu Gln Glu Gly Leu Lys Asn Ala Gly Leu Ala Val
 115 120 125
 Gly Gln Pro Leu Lys Gln Ala Thr Val Gln Met Ile Glu Thr Glu Leu
 130 135 140
 Thr Asn Gln Tyr Ile Ser Gln Gly Tyr Tyr Asn Thr Glu Ile Thr Val
 145 150 155 160
 Lys Gln Thr Met Leu Asp Gly Asn Arg Val Lys Leu Asp Met Thr Phe
 165 170 175
 Ala Glu Gly Lys Pro Ala Arg Val Val Asp Ile Asn Ile Ile Gly Asn
 180 185 190
 Gln His Phe Ser Asp Ala Asp Leu Ile Asp Val Leu Ala Ile Lys Asp
 195 200 205
 Asn Lys Ile Asn Pro Leu Ser Lys Ala Asp Arg Tyr Thr Gln Glu Lys
 210 215 220
 Leu Val Thr Ser Leu Glu Asn Leu Arg Ala Lys Tyr Leu Asn Ala Gly
 225 230 235 240
 Phe Val Arg Phe Glu Ile Lys Asp Ala Lys Leu Asn Ile Asn Glu Asp
 245 250 255
 Lys Asn Arg Ile Phe Val Glu Ile Ser Leu His Glu Gly Glu Gln Tyr
 260 265 270
 Arg Phe Gly Gln Thr Gln Phe Leu Gly Asn Leu Thr Tyr Thr Gln Ala
 275 280 285
 Glu Leu Glu Ala Leu Leu Lys Phe Lys Ala Glu Glu Gly Phe Ser Gln
 290 295 300
 Ala Met Leu Glu Gln Thr Thr Asn Asn Ile Ser Thr Lys Phe Gly Asp
 305 310 315 320
 Asp Gly Tyr Tyr Tyr Ala Gln Ile Arg Pro Val Thr Arg Ile Asn Asp
 325 330 335
 Glu Ser Arg Thr Val Asp Val Glu Tyr Tyr Ile Asp Pro Val His Pro
 340 345 350
 Val Tyr Val Arg Arg Ile Asn Phe Thr Gly Asn Phe Lys Thr Gln Asp
 355 360 365
 Glu Val Leu Arg Arg Glu Met Arg Gln Leu Glu Gly Ala Leu Ala Ser
 370 375 380
 Asn Gln Lys Ile Gln Leu Ser Arg Ala Arg Leu Met Arg Thr Gly Phe
 385 390 395 400
 Phe Lys His Val Thr Val Asp Thr Arg Pro Val Pro Asn Ser Pro Asp
 405 410 415
 Gln Val Asp Val Asn Phe Val Val Glu Gln Pro Ser Gly Ser Ser
 420 425 430
 Thr Ile Ala Ala Gly Tyr Ser Gln Ser Gly Gly Val Thr Phe Gln Phe
 435 440 445
 Asp Val Ser Gln Asn Asn Phe Met Gly Thr Gly Lys His Val Asn Ala
 450 455 460
 Ser Phe Ser Arg Ser Glu Thr Arg Glu Val Tyr Ser Leu Gly Met Thr
 465 470 475 480
 Asn Pro Tyr Phe Thr Val Asn Gly Val Ser Gln Ser Leu Ser Gly Tyr
 485 490 495
 Tyr Arg Lys Thr Lys Tyr Asp Asn Lys Asn Ile Ser Asn Tyr Val Leu
 500 505 510
 Asp Ser Tyr Gly Gly Ser Leu Ser Tyr Gly Tyr Pro Ile Asp Glu Asn
 515 520 525

Gln Arg Ile Ser Phe Gly Leu Asn Ala Asp Asn Thr Lys Leu His Gly
 530 535 540
 Gly Arg Phe Met Gly Ile Ser Asn Val Lys Gln Leu Met Ala Asp Gly
 545 550 555 560
 Gly Lys Ile Gln Val Asp Asn Asn Gly Ile Pro Asp Phe Lys His Asp
 565 570 575
 Tyr Thr Thr Tyr Asn Ala Ile Leu Gly Trp Asn Tyr Ser Ser Leu Asp
 580 585 590
 Arg Pro Val Phe Pro Thr Gln Gly Met Ser His Ser Val Asp Leu Thr
 595 600 605
 Val Gly Phe Gly Asp Lys Thr His Gln Lys Val Val Tyr Gln Gly Asn
 610 615 620
 Ile Tyr Arg Pro Phe Ile Lys Lys Ser Val Leu Arg Gly Tyr Ala Lys
 625 630 635 640
 Leu Gly Tyr Gly Asn Asn Leu Pro Phe Tyr Glu Asn Phe Tyr Ala Gly
 645 650 655
 Gly Tyr Gly Ser Val Arg Gly Tyr Asp Gln Ser Ser Leu Gly Pro Arg
 660 665 670
 Ser Gln Ala Tyr Leu Thr Ala Arg Arg Gly Gln Gln Thr Thr Leu Gly
 675 680 685
 Glu Val Val Gly Gly Asn Ala Leu Ala Thr Phe Gly Ser Glu Leu Ile
 690 695 700
 Leu Pro Leu Pro Phe Lys Gly Asp Trp Ile Asp Gln Val Arg Pro Val
 705 710 715 720
 Ile Phe Ile Glu Gly Gly Gln Val Phe Asp Thr Thr Gly Met Asp Lys
 725 730 735
 Gln Thr Ile Asp Leu Thr Gln Phe Lys Asp Pro Gln Ala Thr Ala Glu
 740 745 750
 Gln Asn Ala Lys Ala Ala Asn Arg Pro Leu Leu Thr Gln Asp Lys Gln
 755 760 765
 Leu Arg Tyr Ser Ala Gly Val Gly Ala Thr Trp Tyr Thr Pro Ile Gly
 770 775 780
 Pro Leu Ser Ile Ser Tyr Ala Lys Pro Leu Asn Lys Lys Gln Asn Asp
 785 790 795 800
 Gln Thr Asp Thr Val Gln Phe Gln Ile Gly Ser Val Phe
 805 810

<210> 5
 <211> 17
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Primer

<400> 5
 actatagggc acgcgtg

17

<210> 6
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Primer

<400> 6
 cctgcgtttg tttgattgag

20

<210> 7
 <211> 61
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide

<400> 7
 aagggcccaa ttacgcagag gggatccaca ggactacagc gagtgaccat tgaaagctta 60
 c 61

<210> 8
 <211> 67
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Oligonucleotide

<400> 8
 aagggcccaa ttacgcagag ggtcgactta ttaaaagaca ctaccaatct ggaactgtac 60
 cgtatcg 67

<210> 9
 <211> 16
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Oligopeptide

<400> 9
 Cys Tyr Ala Lys Pro Leu Asn Lys Lys Gln Asn Asp Gln Thr Asp Thr
 1 5 10 15

<210> 10
 <211> 17
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Oligopeptide

<400> 10
 Tyr Leu Thr Ala Arg Arg Gly Gln Gln Thr Thr Leu Gly Glu Val Val
 1 5 10 15
 Cys

SEQUENCE INFORMATION

BASB027 Polynucleotide and Polypeptide Sequences

SEQ ID NO:1

Moraxella catarrhalis BASB027 polynucleotide sequence from strain ATCC 43617

ATGCGTAATTCATATTTTAAAGGTTTTTCAGGTCAGTGCAATGACAATGGCTGTCATGATG
GTAATGTCAACTCATGCACAAGCGGCGGATTTTATGGCAAATGACATTACCATCACAGGA
CTACAGCGAGTGACCATTGAAAGCTTACAAAGCGTGCTGCCGTTTCGCTTGGGTCAAGTG
GTGAGCGAAAACCAGTTGGCTGATGGTGTCAAAGCACTTTATGCAACAGGCAATTTTTCA
GATGTGCAAGTCTATCATCAAGAAGGGCGTATCATCTATCAGGTAACCGAAAGGCCGTTA
ATCGCTGAGATTAATTTTGGGGCAATCGCTTAATCCAAAAGAAGGTCTACAAGAAGGG
CTAAAAAATGCTGGCTTAGCTGTGGGTCAACCACTAAAACAAGCCACAGTACAGATGATC
GAAACCGAGCTTACCAATCAATATATATCACAAGGCTATTATAATACCGAAATTACTGTC
AAACAGACGATGCTTGATGGTAATCGTGTTAAGCTTGATATGACCTTTGCTGAAGGTAAA
CCTGCACGGGTGGTTGATATTAATATCATTGGCAATCAGCATTTTAGCGATGCAGATTTG
ATTGATGTGCTTGCATTAAAGGATAATAAAATCAATCCACTGTCTAAAGCTGACCGTTAT
ACTCAAGAAAAGCTGGTGACCAGTTTAGAGAATTTGCGTGCTAAATATCTCAATGCAGGG
TTTGTGCGTTTTGAGATTAAAGATGCTAAGCTTAATATTAATGAAGATAAAAACCGTATC
TTTGTGAGATTTTCATTGCATGAAGGTGAGCAATATCGCTTTGGACAGACACAGTTTTTG
GGTAATTTAACTTATACTCAAGCAGAACTTGAGGCACTGCTTAAATTCAAAGCAGAAGAA
GGGTTTTTACAAGCCATGCTTGAGCAAACAACAAACAATATCAGTACCAAATTTGGTGAC
GATGGCTATTATTATGCTCAAATCCGTCCTGTAACACGCATTAATGATGAAAGTCGTACG
GTTGATGTGGAATATTATATTGACCCTGTACACCCTGTCTATGTACGCCGTATTAATTTT
ACAGGTAACCTTTAAGACCCAAGATGAAGTACTCCGTCGTGAGATGCGACAACCTTGAAGGT
GCGTTGGCATCTAATCAAAAAATCCAGCTGTCTCGTGACGCTTGATGCGGACTGGGTTT
TTTAAACATGTTACCGTTGATACTCGTCCAGTACCCAACTCACCTGATCAGGTTGATGTA
AATTTTGTGGTTGAAGAACAACCTTCAGGATCATCAACCATCGCAGCAGGCTACTCTCAA
AGTGGTGGTGTAACTTTTCAATTTGATGTTTCTCAAAATAACTTTATGGGTACAGGTAAG
CACGTCAATGCTTCGTTTTCTCGCTCTGAGACCCGTGAGGTGTATAGTTTGGGTATGACC
AACCCTACTTTTACCGTAAATGGCGTCTCGCAAAGCTTGAGTGGCTACTATCGTAAAACC
AAGTATGATAACAAGAACATTAGTAATTATGTACTTGATTCTTATGGTGGCTCATTAAAGC
TATGGATATCCAATTGATGAAAATCAACGCATAAGCTTTGGTCTGAATGCTGACAATACC
AAGCTTCATGGCGGTCGTTTTATGGGCATTAGTAATGTCAAGCAGCTGATGGCAGATGGT
GGCAAAATTCAAGTGGATAATAATGGCATTCTGATTTTAAGCATGATTACACAACCTAC
AATGCCATTTTGGGGTGAATTATTCAAGTCTAGATCGCCCTGTATTTCCAACCCAAGGC
ATGAGTCATTCTGTAGATTTGACGGTTGGTTTTGGTGATAAACTCATCAAAAAGTGGTT
TATCAAGGCAATATCTATCGCCCATTTATCAAAAAATCAGTCTTGCGTGGATACGCCAAG
TTAGGCTATGGCAATAATTTACCATTTTATGAAAATTTCTATGCAGGCGGCTATGGTTCTG
GTTCTGGCTATGATCAATCCTCTTTGGGTCCACGCTCACAAGCCTATTTGACAGCTCGT
CGTGGTCAACAAACCACACTAGGAGAGGTTGTTGGTGGTAATGCTTTGGCAACTTTCCGGC
AGTGAGCTGATTTTACCTTTGCCATTTAAAGGTGATTGGATAGATCAGGTGCGTCCAGTG
ATATTCATTGAGGGCGGTGAGGTTTTTGATACAACAGGTATGGATAAAACAACCATTTGAT
TTAACCAATTTAAAGACCCACAAGCAACAGCTGAACAAAATGCAAAAGCAGCCAATCGC
CCGCTACTAACCCAAGATAAACAGTTGCGTTATAGTGCTGGTGTGGTGCAACTTGGTAT

ACGCCCATTGGTCCTTTATCTATTAGCTATGCCAAGCCATTGAATAAAAAACAAAATGAT
CAGACCGATACGGTACAGTTCCAGATTGGTAGTGTCTTTTAA

SEQ ID NO:2

Moraxella catarrhalis BASB027 polypeptide sequence deduced from the
polynucleotide sequence of SEQ ID NO:1

MRNSYFKGFQVSAMTMAVMMVMSTHAQAADFMANDITITGLQRVTIESLQSVLPFRLGQV
VSENQLADGVKALYATGNFSDVQVYHQEGRIIYQVTERPLIAEINFEGNRLIPKEGLQEG
LKNAGLAVGQPLKQATVQMIETELTNQYISQGYNTEITVKQTMLDGNRVKLDMTFAEGK
PARVVDINIIGNQHFSADLIDVLAIKDNKINPLSKADRYTQEKLVTSLLENLRKAYLNAG
FVRFEIKDAKLNINEDKNRIFVEISLHEGEQYRFGQTQFLGNLTYTQAELEALLKFKAEE
GFSQAMLEQTTNNISTKFGDDGYYYAQIRPVTRINDESRTVDVEYYIDPVHPVYVRRINF
TGNFKTQDEVLRREMRQLEGALASNQKIQLSRARLMRTGFFKHVTVDTRPVPNSPDQVDV
NFWVEEQPSGSSTIAAGYSQSGGVTFQFDVSQNNFMGTGKHVNASFRRSETREVYSLGMT
NPYFTVNGVSQSLSGYYRKTKYDNKNISNYVLDSYGGSLSYGYPIDENQRISFGLNADNT
KLHGGRFMGISNVKQLMADGGKIQVDNNGIPDFKHDYTTYNAILGWNYSLLDRPVFPPTQG
MSHSVLDLTVFGDKTHQKVYVYQGNIRPFIKKSVLRGYAKLGYGNNLPFYENFYAGGYGS
VRGYDQSSSLGPRSQAAYLTARRGQQTTLGEVVGGNALATFGSELILPLPFKGDWIDQVRPV
IFIEGGQVFDTTGMDKQTDIDLTQFKDPQATAEQNAKAANRPLLTDQDKQLRYSAGVGATWY
TPIGPLSISYAKPLNKKQNDQTDQVQFQIGSVF

SEQ ID NO:3

Moraxella catarrhalis BASB027 polynucleotide sequence from strain ATCC 43617

ATGCGTAATTCATATTTTAAAGGTTTTTCAGGTCAGTGCAATGACAATGGCTGTCATGATG
GTAATGTCAACTCATGCACAAGCGGCGGATTTTATGGCAAATGACATTACCATCACAGGA
CTACAGCGAGTGACCATTGAAAGCTTACAAAGCGTGCTGCCGTTTCGCTTGGGTCAAGTG
GTGAGCGAAAACAGTTGGCTGATGGTGTCAAAGCACTTTATGCAACAGGCAATTTTCA
GATGTGCAAGTCTATCATCAAGAAGGGCGTATCATCTATCAGGTAACCGAAAGGCCGTTA
ATCGCTGAGATTAATTTTGAGGGCAATCGCTTAATTCCAAAAGAAGGTCTACAAGAAGGG
CTAAAAAATGCTGGCTTAGCTGTGGGTCAACCACTAAAACAAGCCACAGTACAGATGATC
GAAACCGAGCTTACCAATCAATATATATCACAAGGCTATTATAATACCGAAATTACTGTC
AAACAGACGATGCTTGATGGTAATCGTGTTAAGCTTGATATGACCTTTGCTGAAGGTAAA
CCTGCACGGGTGGTTGATATTAATATCATTGGCAATCAGCATTTTAGCGATGCAGATTTG
ATTGATGTGCTTGCATTAAGGATAATAAAATCAATCCACTGTCTAAAGCTGACCGTTAT
ACTCAAGAAAAGCTGGTGACCAGTTTAGAGAATTTGCGTGCTAAATATCTCAATGCAGGG
TTTGTGCGTTTTGAGATTAAAGATGCTAAGCTTAATATTAATGAAGATAAAAACCGTATC
TTTGTGAGATTTTATTGCATGAAGGTGAGCAATATCGCTTTGGACAGACACAGTTTTTG
GGTAATTTAACTTATACTCAAGCAGAACTTGAGGCACTGCTTAAATTCAAAGCAGAAGAA

GGGTTTTTCACAAGCCATGCTTGAGCAAACAACAACAATATCAGTACCAAATTTGGTGAC
 GATGGCTATTATTATGCTCAAATCCGTCCTGTAACACGCATTAATGATGAAAGTCGTACG
 GTTGATGTGGAATATTATATTGACCCTGTACACCCTGTCTATGTACGCCGTATTAATTTT
 ACAGGTAACCTTTAAGACCCAAGATGAAGTACTCCGTCGTGAGATGCGACAACCTGAAGGT
 GCGTTGGCATCTAATCAAAAAATCCAGCTGTCTCGTGACGCTTGATGCGGACTGGGTTT
 TTAAACATGTTACCGTTGATACTCGTCCAGTACCCAACCTCACCTGATCAGGTTGATGTA
 AATTTTGTGGTTGAAGAACAACCTTCAGGATCATCAACCATCGCAGCAGGCTACTCTCAA
 AGTGGTGGTGTAACCTTTCAATTTGATGTTTCTCAAATAACTTTATGGGTACAGGTAAG
 CACGTCAATGCTTCGTTTTCTCGCTCTGAGACCCGTGAGGTGTATAGTTTGGGTATGACC
 AACCCTACTTTACCGTAAATGGCGTCTCGCAAAGCTTGAGTGGCTACTATCGTAAACC
 AAGTATGATAACAAGAACATTAGTAATTATGTACTTGATTCTTATGGTGGCTCATTAGC
 TATGGATATCCAATTGATGAAAATCAACGCATAAGCTTTGGTCTGAATGCTGACAATACC
 AAGCTTCATGGCGGTGCTTTTTATGGGCATTAGTAATGTCAAGCAGCTGATGGCAGATGGT
 GGCAAATTCAGTGGATAATAATGGCATTCTGATTTTAAGCATGATTACACAACCTAC
 AATGCCATTTTGGGGTGAATTAATCAAGTCTAGATCGCCCTGTATTTCCAACCCAAGGC
 ATGAGTCATTCTGTAGATTTGACGGTTGGTTTTGGTGATAAACTCATCAAAAGTGGTT
 TATCAAGGCAATATCTATCGCCATTTATCAAAAAATCAGTCTTGCGTGGATACGCCAAG
 TTAGGCTATGGCAATAATTTACCATTTTATGAAAATTTCTATGCAGGCGGCTATGGTTCG
 GTTCGTGGCTATGATCAATCCTCTTTGGGTCCACGCTCACAAGCCTATTTGACAGCTCGT
 CGTGGTCAACAAACCACACTAGGAGAGGTTGTTGGTGGTAATGCTTTGGCAACTTTCGGC
 AGTGAGCTGATTTTACCTTTGCCATTTAAAGGTGATTGGATAGATCAGGTGCGTCCAGTG
 ATATTCATTGAGGGCGGTCAGGTTTTTGATACAACAGGTATGGATAAACAACCATTTGAT
 TTAACCCAATTTAAAGACCCACAAGCAACAGCTGAACAAAATGCAAAAGCAGCCAATCGC
 CCGCTACTAACCCAAGATAAACAGTTGCGTTATAGTGCTGGTGTGGTGCAACTTGGTAT
 ACGCCCATTTGGTCTTTATCTATTAGCTATGCCAAGCCATTGAATAAAAAACAAATGAT
 CAGACCGATACGGTACAGTTCAGATTGGTAGTGTCTTTTAA

SEQ ID NO:4

Moraxella catarrhalis BASB027 polypeptide sequence deduced from the
 polynucleotide sequence of SEQ ID NO:3

MRNSYFKGFQVSAMTMAVMMVMSTHAQAADFMANDITITGLQRVTIESLQSVLPFRLGQV
 VSENQLADGVKALYATGNFSDVQVYHQEGRIIYQVTERPLIAEINFEGNRLIPKEGLQEG
 LKNAGLAVGQPLKQATVQMIETELTNQYISQGYNTEITVKQTMLDGNRVKLDMTFAEGK
 PARVVDINIIGNQHFSADLIDVLAIKDNKINPLSKADRYTQEKLVTSLLENLRAKYLNAG
 FVRFEIKDAKLNINEDKNRIFVEISLHEGEQYRFGQTQFLGNLTYTQAELEALLKFKAEE
 GFSQAMLEQTTNNISTKFGDDGYYYAQIRPVTRINDESRTVDVEYYIDPVHPVYVRRINF
 TGNFKTQDEVLRREMRQLEGALASNQKIQLSRARLMRTGFFKHVTVDTRPVPNSPDQVDV
 NFVVEEQPSGSSTIAAGYSQSGGVTFQFDVSNQNFMTGKHVNASFSRSETREVYSLGMT
 NPYFTVNGVSQSLSGYYRKYDNKNISNYVLDSYGGSLSYGYPIDENQRISFGLNADNT
 KLHGGRFMGISNVKQLMADGGKIQVDNNGIPDFKHDTTYNAILGWNYSLLDRPVFPTQG
 MSHSVDLTVGFQDKTHQKVYQGNIRPFIKSVLRGYAKLGYGNNLPFYENFYAGGYGS
 VRGYDQSSLGPRSQAYLTARRGQQTTLGEVVGGNALATFGSELILPLPKGDWIDQVRPV
 IFIEGGQVFDTTGMDKQTDILTQFKDPQATAEQNAKAANRPLLTDQKQLRYSAGVGATWY

TPIGPLSISYAKPLNKKQNDQTDTVQFQIGSVF

SEQ ID NO:5

ACT ATA GGG CAC GCG TG

SEQ ID NO:6

CCT GCG TTT GTT TGA TTG AG

SEQ ID NO:7

AAG GGC CCA ATT ACG CAG AGG GGA TCC ACA GGA CTA CAG CGA GTG
ACC ATT GAA AGC TTA C

SEQ ID NO:8

AAG GGC CCA ATT ACG CAG AGG GTC GAC TTA TTA AAA GAC ACT ACC
AAT CTG GAA CTG TAC CGT ATC G

SEQ ID NO:9

CYAKPLNKKQNDQTD

SEQ ID NO:10

YLTARRGQQTTLGEVVC

DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

**BASB027 PROTEINS AND GENES FROM MORAXELLA CATARRHALIS ,
ANTIGENS, ANTIBODIES, AND USES**

the specification of which (check one)

☐ is attached hereto.

☒ was filed on 31 May 1999

as Serial No. PCT/EP99/03822

and was amended on

(if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the patentability as defined in Title 37, Code of Federal Regulations, Section 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, Section 119(a)-(d) or Section 365(b) of any foreign application(s) for patent or inventor's certificate, or Section 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below any foreign application for patent or Inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Number	Country	Filing Date	Priority Claimed
9811945.6	Great Britain	03 June 1998	Yes

I hereby claim the benefit under Title 35, United States Code, Section 119(e) of any United States provisional application(s) listed below.

Application Number Filing Date

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States application(s) or Section 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

Serial No.	Filing Date	Status
------------	-------------	--------


I hereby appoint the practitioners associated with the Customer Number provided below to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith, and direct that all correspondence be addressed to that Customer Number:

Customer Number 25,308

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge

that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full Name of Inventor: Carlota VINALS y de BASSOLS

Inventor's Signature: 

Date: 15.02.01


Residence: Brussels, Belgium

BEX

Citizenship: Belgian

Post Office Address: SmithKline Beecham Corporation
Corporate Intellectual Property - UW2220
P.O. Box 1539
King of Prussia, Pennsylvania 19406-0939

United States Patent & Trademark Office
Office of Initial Patent Examination — Scanning Division



Application deficiencies found during scanning:

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☒ *Scanned copy is best available. drawings are dark*